

Product datasheet for TP303667L

Cirhin (UTP4) (NM_032830) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human cirrhosis, autosomal recessive 1A (cirhin) (CIRH1A), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC203667 protein sequence Red=Cloning site Green=Tags(s)

MGEFKVHRVRRFFNYVPSGIRCVAYNNQSNRLAVSRTDGTVEIYNLSANYFQEKFFPGHESRATEALCWAE
GQRLFSAGLNGEIMEYDLQALNIKYAMDAFGGPIWSMAASPSGSQLLVGCEDGSVKLFQITPDKIQFERN
FDRQKSRILSLSWHPSGTHIAAGSIDYISVFDVKSGSAVHKMIVDRQYMGVSKRKCIVWGVAFVLSGTII
SVDSAGKVQFWDSATGTLVKSHLIANADVQSIADVADQEDSFVVGTAEGTVFHFQLVPVTSNSSEKQWVRT
KPFQHHTHDVVRTVAHSPTALISGGTDTHLVFRPLMEKVEVKNYDAALRKITFPHRCLISCSKRRQLLLFQ
FAHHELELWRLGSTVATGKNGDTLPLSKNADHLLHLKTKGPENIICSCISPCGSWIAYSTVSRFFLYRLNY
EHDNISLKRVSMPAFLRSALQILFSEDSTKLFVASNQGALHIVQLSGGSFKHLHAFQPQSGTVEAMCLL
AVSPDGNWLAASGTSAGVHVYVNVKQLKLHCTVPAYNFPVTAMAIAPNTNNLVIAHSDQQVF EYSIPDKQY
TDWSRTVQKQGFHHLWLQRDTPITHISFHPKRPMHILLHDAYMFCIIDKSLPLPNDKTLNPNFPPTNES
DVIRRRATAHAFKISKIYKPLLFMDLLDERTLVAVERPLDDIIAQLPPPIK KKKKFGT

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	76.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_116219
Locus ID:	84916
UniProt ID:	Q969X6
RefSeq Size:	2268
Cytogenetics:	16q22.1
RefSeq ORF:	2058
Synonyms:	CIRH1A; CIRHIN; NAIC; TEX292
Summary:	This gene encodes a WD40-repeat-containing protein that is localized to the nucleolus. Mutation of this gene causes North American Indian childhood cirrhosis, a severe intrahepatic cholestasis that results in transient neonatal jaundice, and progresses to periportal fibrosis and cirrhosis in childhood and adolescence. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2016]

Product images:



Coomassie blue staining of purified UTP4 protein (Cat# [TP303667]). The protein was produced from HEK293T cells transfected with UTP4 cDNA clone (Cat# [RC203667]) using MegaTran 2.0 (Cat# [TT210002]).