

## Product datasheet for TP303699L

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Macro H2A.2 (H2AFY2) (NM\_018649) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human H2A histone family, member Y2 (H2AFY2), 1 mg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC203699 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MSGRSGKKKMSKLSRSARAGVIFPVGRLMRYLKKGTFKYRISVGAPVYMAAVIEYLAAEILELAGNAARD NKKARIAPRHILLAVANDEELNQLLKGVTIASGGVLPRIHPELLAKKRGTKGKSETILSPPPEKRGRKAT SGKKGGKKSKAAKPRTSKKSKPKDSDKEGTSNSTSEDGPGDGFTILSSKSLVLGQKLSLTQSDISHIGSM RVEGIVHPTTAEIDLKEDIGKALEKAGGKEFLETVKELRKSQGPLEVAEAAVSQSSGLAAKFVIHCHIPQ WGSDKCEEQLEETIKNCLSAAEDKKLKSVAFPPFPSGRNCFPKQTAAQVTLKAISAHFDDSSASSLKNVY

**FLLFDSESIGIYVQEMAKLDAK** 

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK

**Predicted MW:** 39.9 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

**Storage:** Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** <u>NP 061119</u>

**Locus ID:** 55506





Synonyms:

UniProt ID: <u>Q9P0M6</u>, <u>A0A024QZP6</u>

RefSeq Size: 2181

Cytogenetics: 10q22.1 RefSeq ORF: 1116

**Summary:** Histones are basic nuclear proteins that are responsible for the nucleosome structure of the


chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent histone that is a member of the histone H2A family. It replaces conventional H2A histones in a subset of nucleosomes where it represses transcription and may participate in stable X chromosome inactivation. [provided

by RefSeq, Oct 2015]

**Protein Pathways:** Systemic lupus erythematosus

H2AFY2

## **Product images:**



Coomassie blue staining of purified H2AFY2 protein (Cat# [TP303699]). The protein was produced from HEK293T cells transfected with H2AFY2 cDNA clone (Cat# [RC203699]) using MegaTran 2.0 (Cat# [TT210002]).