

Product datasheet for TP303773

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TFAP4 (NM 003223) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human transcription factor AP-4 (activating enhancer binding protein

4) (TFAP4), 20 μg

Species: Human Expression Host: HEK293T

Expression cDNA Clone >RC203773 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MEYFMVPTQKVPSLQHFRKTEKEVIGGLCSLANIPLTPETQRDQERRIRREIANSNERRRMQSINAGFQS LKTLIPHTDGEKLSKAAILQQTAEYIFSLEQEKTRLLQQNTQLKRFIQELSGSSPKRRRAEDKDEGIGSP DIWEDEKAEDLRREMIELRQQLDKERSVRMMLEEQVRSLEAHMYPEKLKVIAQQVQLQQQQEQVRLLHQE KLEREQQQLRTQLLPPPAPTHHPTVIVPAPPPPPSHHINVVTMGPSSVINSVSTSRQNLDTIVQAIQHIE GTQEKQELEEEQRRAVIVKPVRSCPEAPTSDTASDSEASDSDAMDQSREEPSGDGELP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 38.5 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 003214

Locus ID: 7023



UniProt ID: Q01664

RefSeq Size: 2147

Cytogenetics: 16p13.3 RefSeq ORF: 1014

Synonyms: AP-4; bHLHc41


Summary: Transcription factors of the basic helix-loop-helix-zipper (bHLH-ZIP) family contain a basic

domain, which is used for DNA binding, and HLH and ZIP domains, which are used for oligomerization. Transcription factor AP4 activates both viral and cellular genes by binding to the symmetrical DNA sequence CAGCTG (Mermod et al., 1988 [PubMed 2833704]; Hu et al.,

1990 [PubMed 2123466]).[supplied by OMIM, Jul 2009]

Protein Families: Druggable Genome, Transcription Factors

Product images:



Coomassie blue staining of purified TFAP4 protein (Cat# TP303773). The protein was produced from HEK293T cells transfected with TFAP4 cDNA clone (Cat# [RC203773]) using MegaTran 2.0 (Cat# [TT210002]).