

# **Product datasheet for TP303773M**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### TFAP4 (NM\_003223) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human transcription factor AP-4 (activating enhancer binding protein

4) (TFAP4), 100 μg

Species: Human Expression Host: HEK293T

**Expression cDNA Clone** >RC203773 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MEYFMVPTQKVPSLQHFRKTEKEVIGGLCSLANIPLTPETQRDQERRIRREIANSNERRRMQSINAGFQS LKTLIPHTDGEKLSKAAILQQTAEYIFSLEQEKTRLLQQNTQLKRFIQELSGSSPKRRRAEDKDEGIGSP DIWEDEKAEDLRREMIELRQQLDKERSVRMMLEEQVRSLEAHMYPEKLKVIAQQVQLQQQQEQVRLLHQE KLEREQQQLRTQLLPPPAPTHHPTVIVPAPPPPPPSHHINVVTMGPSSVINSVSTSRQNLDTIVQAIQHIE GTQEKQELEEEQRRAVIVKPVRSCPEAPTSDTASDSEASDSDAMDQSREEPSGDGELP

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK

Predicted MW: 38.5 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

**Storage:** Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 003214

**Locus ID:** 7023





RefSeq ORF:

#### TFAP4 (NM\_003223) Human Recombinant Protein - TP303773M

**UniProt ID:** Q01664

RefSeq Size: 2147

Cytogenetics: 16p13.3 1014

Synonyms: AP-4; bHLHc41


**Summary:** Transcription factors of the basic helix-loop-helix-zipper (bHLH-ZIP) family contain a basic

> domain, which is used for DNA binding, and HLH and ZIP domains, which are used for oligomerization. Transcription factor AP4 activates both viral and cellular genes by binding to the symmetrical DNA sequence CAGCTG (Mermod et al., 1988 [PubMed 2833704]; Hu et al.,

1990 [PubMed 2123466]).[supplied by OMIM, Jul 2009]

**Protein Families:** Druggable Genome, Transcription Factors

## **Product images:**



Coomassie blue staining of purified TFAP4 protein (Cat# [TP303773]). The protein was produced from HEK293T cells transfected with TFAP4 cDNA clone (Cat# [RC203773]) using MegaTran 2.0

(Cat# [TT210002]).