

Product datasheet for TP303788L

PSTPIP1 (NM_003978) Human Recombinant Protein


Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human proline-serine-threonine phosphatase interacting protein 1 (PSTPIP1), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC203788 protein sequence Red =Cloning site Green =Tags(s) MMPQLQFKDAFWCRDFTAHTGYEVLLQRLLDGRKMCKDMEELLRQRAQAEERYGKELVQIARKAGGQT EI NSLRASFDLQKQMMENVGSSHIQLALTREELRSLEEFRRQKEQRKKYEAVMDRVQSKLSLYKKAMES KKTYEQKCRDADDAEQAFERISANGHQKQVEKSNKARQCKDSATEAERVYRQSIQLEKVRAEWEQEH R TTCEAFQLQEFDRILTILRNALWVHSNQLSMQCVKDDELYEEVRLTLEGCSIDADIDSFIQAKSTGTEPPA PVPYQNYDREVPLTSSPGIQPSCGMIKRFSGLLHGSPKTTSLAASAATETLTPTPERNEGVYTAIAV QEIQGNPASPAQEYRALYDYTAQNPDELDSLQAGDILEVILEGEDGWWTVERNQGQGFVPGSYLEKL TR TRPLE QKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	47.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.


[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_003969
Locus ID:	9051
UniProt ID:	O43586
RefSeq Size:	1870
Cytogenetics:	15q24.3
RefSeq ORF:	1248
Synonyms:	CD2BP1; CD2BP1L; CD2BP1S; H-PIP; PAPAS; PSTPIP
Summary:	This gene encodes a cytoskeletal protein that is highly expressed in hemopoietic tissues. This protein functions via its interaction with several different proteins involved in cytoskeletal organization and inflammatory processes. It binds to the cytoplasmic tail of CD2, an effector of T cell activation and adhesion, downregulating CD2-triggered adhesion. It binds PEST-type protein tyrosine phosphatases (PTP) and directs them to c-Abl kinase to mediate c-Abl dephosphorylation, thereby, regulating c-Abl activity. It also interacts with pyrin, which is found in association with the cytoskeleton in myeloid/monocytic cells and modulates immunoregulatory functions. Mutations in this gene are associated with PAPA (pyogenic sterile arthritis, pyoderma gangrenosum, and acne) syndrome. It is hypothesized that the disease-causing mutations compromise physiologic signaling necessary for the maintenance of a proper inflammatory response. [provided by RefSeq, Mar 2016]
Protein Families:	Druggable Genome
Protein Pathways:	NOD-like receptor signaling pathway

Product images:



Coomassie blue staining of purified PSTPIP1 protein (Cat# [TP303788]). The protein was produced from HEK293T cells transfected with PSTPIP1 cDNA clone (Cat# [RC203788]) using MegaTran 2.0 (Cat# [TT210002]).