

## Product datasheet for TP303795L

### CCR4 NOT transcription complex subunit 3 (CNOT3) (NM\_014516) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human CCR4-NOT transcription complex, subunit 3 (CNOT3), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC203795 protein sequence Red=Cloning site Green=Tags(s)

MADKRKLQGEIDRCLKKVEGVEQFEDIWQKLHNAANANQKEKYEADLKKKEIKKLQRLRDQIKTWASNE  
IKDKRQLIDNRKLIETQMERFKVVERETKTAKYSKEGLGLAQKVDPAQKEKEEVGQWLTNTIDTLNMQVD  
QFESEVESLSVQTRKKKGDKDKQDRIEGLKRHIEKHRYHVRMLETILRMLDNDNSILVDAIRKIKDDVEYY  
VDSSQDPDFEENEFLYDDLLEDIPQALVATSPSSHMEDEFNQSSSTPTSTTSSSPIPPSPANCTTE  
NSEDKKRGRSTDSEVSQSPAKNGSKPVHSNQHPQSPAVPPTYPSGPPPAASALSTTPGNNGVPAPAAPP  
SALGPKASPAPSHNSGTPAPYAQAVAPPAPSGPSTTQPRPPSVQPSGGGGGGGGSSSSSSNSAGGGA  
GKQNGATSYSSVADSPAVALSSSGGNNASSQALGPPSGPHNPPPSTSKEPSAAAPTGAGGVAPGSGNN  
SGGPSLLVPLPVNPPSPTPSFSDAKAAGALLNGPPQFSTAPEIKAPEPLSSLKSMASRAAIISSGIEDPV  
PTLHLTERDIILSSTSAPPASAQPPLQLSEVNIPLSLGVCPLGPVPLTKEQLYQQAMEEAAWHHMPHPSD  
SERIRQYLPRNPCPTPPYHHQMPPPHSDTVEFYQRLSTETLFFIFYLEGTKAQYLAALKKQSWRFHT  
KYMMWFQRHEEPTITDEFEQGTIYFDYEKWGQRKKEGFTFEYRYLEDRLQ


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	81.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.



[View online »](#)

<b>Storage:</b>	Store at -80°C.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_055331</a>
<b>Locus ID:</b>	4849
<b>UniProt ID:</b>	<a href="#">O75175</a> , <a href="#">A0A024R4R3</a>
<b>RefSeq Size:</b>	2908
<b>Cytogenetics:</b>	19q13.42
<b>RefSeq ORF:</b>	2259
<b>Synonyms:</b>	IDDSADF; LENG2; NOT3; NOT3H
<b>Summary:</b>	Component of the CCR4-NOT complex which is one of the major cellular mRNA deadenylases and is linked to various cellular processes including bulk mRNA degradation, miRNA-mediated repression, translational repression during translational initiation and general transcription regulation. Additional complex functions may be a consequence of its influence on mRNA expression. May be involved in metabolic regulation; may be involved in recruitment of the CCR4-NOT complex to deadenylation target mRNAs involved in energy metabolism. Involved in mitotic progression and regulation of the spindle assembly checkpoint by regulating the stability of MAD1L1 mRNA. Can repress transcription and may link the CCR4-NOT complex to transcriptional regulation; the repressive function may involve histone deacetylases. Involved in the maintenance of embryonic stem (ES) cell identity.[UniProtKB/Swiss-Prot Function]
<b>Protein Families:</b>	Transcription Factors
<b>Protein Pathways:</b>	RNA degradation

**Product images:**

Coomassie blue staining of purified CNOT3 protein (Cat# [TP303795]). The protein was produced from HEK293T cells transfected with CNOT3 cDNA clone (Cat# [RC203795]) using MegaTran 2.0 (Cat# [TT210002]).