

## Product datasheet for **TP303877**

### **MOSC2 (MARC2) (NM\_017898) Human Recombinant Protein**

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human MOCO sulphurase C-terminal domain containing 2 (MOSC2), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC203877 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MGASSSSALARLGLPARPWPRWLGVAALGLAAVALGTVAWRRRAWPRRRRRLQQVGTVAKLWIYPVKCKG  
VPVSEAECTAMGLRSGNLRDRFWLVIKEDGHMVTARQEPRLVLISIIYENNCLIFRAPDMDQLVLPKQP  
SSNKLHNCRIFGLDIKGRDCGNEAAKWFTNFKTEAYRLVQFETNMKGRTSRKLLPTLDQNFQVAYPDYC  
PLLIMTDASLVLDLNRMEKKMKMENFRPNIVVTGCDAFEEDTWEDELLIGSVEVKKVMACPRCILTTPDP  
TGVIDRKQPLDTLKSRYRLCDPSERELYKLSPLFGIYYSVEKIGSLRVGDPVYRMV

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

Tag:	C-Myc/DDK
Predicted MW:	37.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u><a href="#">NP_060368</a></u>
Locus ID:	54996



[View online »](#)

UniProt ID: [Q969Z3](#)

RefSeq Size: 1618

Cytogenetics: 1q41


RefSeq ORF: 1005

Synonyms: MARC2; MOSC2

**Summary:** The protein encoded by this gene is an enzyme found in the outer mitochondrial membrane that reduces N-hydroxylated substrates. The encoded protein uses molybdenum as a cofactor and cytochrome b5 type B and NADH cytochrome b5 reductase as accessory proteins. One type of substrate used is N-hydroxylated nucleotide base analogues, which can be toxic to a cell. Other substrates include N(omega)-hydroxy-L-arginine (NOHA) and amidoxime prodrugs, which are activated by the encoded enzyme. Multiple transcript variants encoding the different isoforms have been found for this gene. [provided by RefSeq, Sep 2016]

**Protein Families:** Transmembrane

### Product images:



Coomassie blue staining of purified MARC2 protein (Cat# TP303877). The protein was produced from HEK293T cells transfected with MARC2 cDNA clone (Cat# [RC203877]) using MegaTran 2.0 (Cat# [TT210002]).