

Product datasheet for **TP303938L**

LITAF (NM_004862) Human Recombinant Protein


Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human lipopolysaccharide-induced TNF factor (LITAF), transcript variant 1, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC203938 protein sequence Red =Cloning site Green =Tags(s) MSVPGPYQAATGPSSAPSAPPSYEETVAVNSYYPTPPAPMPGPTTGLVTGPDGKGMNPPSYTQPAPIPN NNPITVQTVYVQHPITFLDRPIQMCCPSCNKMIVSQLSYNAGALTWLSGSLCLLGCIAGCCFIPFCVDA LQDVDHYCPNCRALLGTYKRL TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	16.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_004853</u>
Locus ID:	9516
UniProt ID:	<u>Q99732</u>


[View online »](#)

RefSeq Size:	2642
Cytogenetics:	16p13.13
RefSeq ORF:	483
Synonyms:	PIG7; SIMPLE; TP53I7
Summary:	Lipopolysaccharide is a potent stimulator of monocytes and macrophages, causing secretion of tumor necrosis factor-alpha (TNF-alpha) and other inflammatory mediators. This gene encodes lipopolysaccharide-induced TNF-alpha factor, which is a DNA-binding protein and can mediate the TNF-alpha expression by direct binding to the promoter region of the TNF-alpha gene. The transcription of this gene is induced by tumor suppressor p53 and has been implicated in the p53-induced apoptotic pathway. Mutations in this gene cause Charcot-Marie-Tooth disease type 1C (CMT1C) and may be involved in the carcinogenesis of extramammary Paget's disease (EMPD). Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Dec 2014]
Protein Families:	Druggable Genome, Transcription Factors

Product images:



Coomassie blue staining of purified LITAF protein (Cat# [TP303938]). The protein was produced from HEK293T cells transfected with LITAF cDNA clone (Cat# [RC203938]) using MegaTran 2.0 (Cat# [TT210002]).