

Product datasheet for TP303944L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MAX (NM_145114) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human MYC associated factor X (MAX), transcript variant 4, 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC203944 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MSDNDDIEVESDEEQPRFQSAADKRAHHNALERKRRDHIKDSFHSLRDSVPSLQGEKLYFLFWKLCTPVL

HRQSLMQKCHTFISSYQVHKKKECKI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 11.3 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 660089

 Locus ID:
 4149

 UniProt ID:
 P61244

 RefSeq Size:
 937

Cytogenetics: 14q23.3



MAX (NM_145114) Human Recombinant Protein - TP303944L

RefSeq ORF: 288

Synonyms: bHLHd4

Summary: The protein encoded by this gene is a member of the basic helix-loop-helix leucine zipper


(bHLHZ) family of transcription factors. It is able to form homodimers and heterodimers with other family members, which include Mad, Mxi1 and Myc. Myc is an oncoprotein implicated in cell proliferation, differentiation and apoptosis. The homodimers and heterodimers compete for a common DNA target site (the E box) and rearrangement among these dimer forms provides a complex system of transcriptional regulation. Mutations of this gene have been reported to be associated with hereditary pheochromocytoma. A pseudogene of this gene is located on the long arm of chromosome 7. Alternative splicing results in multiple transcript

variants. [provided by RefSeq, Aug 2012]

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: MAPK signaling pathway, Pathways in cancer, Small cell lung cancer

Product images:



Coomassie blue staining of purified MAX protein (Cat# [TP303944]). The protein was produced from HEK293T cells transfected with MAX cDNA clone (Cat# [RC203944]) using MegaTran 2.0 (Cat# [TT210002]).