

## Product datasheet for TP304031

### NC2 alpha (DRAP1) (NM\_006442) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human DR1-associated protein 1 (negative cofactor 2 alpha) (DRAP1), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC204031 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)  MPSKKKKYNARFPFARIKKIMQTDEEIGKVA AAVPVIISRALELFLESLLKKACQVTQSRNAKTMTTSHL KQCIELEQQFDLFLKDLVASVPDMQGDGEDNHMDGDKGARRGRKPGSGGRKNGGMGTSKDKKLSGTDSEQ EDESEDTDGEEETSQPPPQASHPSAHFQSPPTPLPFAS TLPLPPAPPGPSAPDEEDEEDYDS  <b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
Tag:	C-Myc/DDK
Predicted MW:	22.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_006433</a>
Locus ID:	10589
UniProt ID:	<a href="#">Q14919</a>



[View online »](#)

RefSeq Size: 1022

Cytogenetics: 11q13.1


RefSeq ORF: 615

Synonyms: NC2-alpha

**Summary:** Transcriptional repression is a general mechanism for regulating transcriptional initiation in organisms ranging from yeast to humans. Accurate initiation of transcription from eukaryotic protein-encoding genes requires the assembly of a large multiprotein complex consisting of RNA polymerase II and general transcription factors such as TFIIA, TFIIB, and TFIID. DR1 is a repressor that interacts with the TATA-binding protein (TBP) of TFIID and prevents the formation of an active transcription complex by precluding the entry of TFIIA and/or TFIIB into the preinitiation complex. The protein encoded by this gene is a corepressor of transcription that interacts with DR1 to enhance DR1-mediated repression. The interaction between this corepressor and DR1 is required for corepressor function and appears to stabilize the TBP-DR1-DNA complex. [provided by RefSeq, Jul 2008]

**Protein Families:** Transcription Factors

### Product images:



Coomassie blue staining of purified DRAP1 protein (Cat# TP304031). The protein was produced from HEK293T cells transfected with DRAP1 cDNA clone (Cat# [RC204031]) using MegaTran 2.0 (Cat# [TT210002]).