

Product datasheet for TP304056L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD99 (NM_002414) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Homo sapiens CD99 molecule (CD99), transcript variant 1, 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC204056 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MARGAALALLLFGLLGVLVAAPDGGFDLSDALPDNENKKPTAIPKKPSAGDDFDLGDAVVDGENDDPRPP NPPKPMPNPNHPSSSGSFSDADLADGVSGGEGKGGSDGGGSHRKEGEEADAPGVIPGIVGAVVVAVAG

AISSFIAYQKKKLCFKENAEQGEVDMESHRNANAEPAVQRTLLEK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 16.7 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling

conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 002405

 Locus ID:
 4267

 UniProt ID:
 P14209

 RefSeq Size:
 1255



Cytogenetics: X;Y

RefSeq ORF: 555

Synonyms: HBA71; MIC2; MIC2X; MIC2Y; MSK5X

Summary: The protein encoded by this gene is a cell surface glycoprotein involved in leukocyte migration,

T-cell adhesion, ganglioside GM1 and transmembrane protein transport, and T-cell death by a caspase-independent pathway. In addition, the encoded protein may have the ability to rearrange the actin cytoskeleton and may also act as an oncosuppressor in osteosarcoma. This

gene is found in the pseudoautosomal region of chromosomes X and Y and escapes X-


chromosome inactivation. There is a related pseudogene located immediately adjacent to this

locus. [provided by RefSeq, Mar 2016]

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs), Leukocyte transendothelial migration

Product images:



Coomassie blue staining of purified CD99 protein (Cat# [TP304056]). The protein was produced from HEK293T cells transfected with CD99 cDNA clone (Cat# [RC204056]) using MegaTran 2.0 (Cat# [TT210002]).