

Product datasheet for **TP304138**

TAF10 (NM_006284) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins
Description: Recombinant protein of human TAF10 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 30kDa (TAF10), 20 µg

Species: Human

Expression Host: HEK293T

Expression cDNA Clone or AA Sequence: >RC204138 protein sequence
Red=Cloning site **Green**=Tags(s)

MSCSGSGADPEAAPASAASAPGPAPPVSAPAALPSSTAAENKASPAGTAGGPGAGAAAGGTGPLAARAGE
PAERRGAAPVSAGGAAPPEGAIISNGVYVLPAAANGDVKPVVSTPLVDFLMQLEDYTPTIPDAVTGYLNL
RAGFEASDPRIIRLISLAAQKFISDIANDALQHCKMKGTASGSSRSKSKDRKYTLTMEDLTPALSEYGIN
VKKPHYFT

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 21.5 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_006275](#)

Locus ID: 6881



[View online »](#)


UniProt ID: [Q12962](#)
RefSeq Size: 834
Cytogenetics: 11p15.4
RefSeq ORF: 654
Synonyms: TAF2A; TAF2H; TAFII30

Summary: Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes one of the small subunits of TFIID that is associated with a subset of TFIID complexes. Studies with human and mammalian cells have shown that this subunit is required for transcriptional activation by the estrogen receptor, for progression through the cell cycle, and may also be required for certain cellular differentiation programs. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Basal transcription factors

Product images:



Coomassie blue staining of purified TAF10 protein (Cat# TP304138). The protein was produced from HEK293T cells transfected with TAF10 cDNA clone (Cat# [RC204138]) using MegaTran 2.0 (Cat# [TT210002]).