

Product datasheet for TP304139

RGS14 (NM_006480) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human regulator of G-protein signaling 14 (RGS14), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC204139 protein sequence Red=Cloning site Green=Tags(s)

MPGKPKHLGVPNGRMVLAVSDGELSSTTGPQQGEGRGSSLSIHSPLSGPSSPFPTEEQPVASWALSFER
LLQDPLGLAYFTEFLKKEFSAENVTFWKACERFQQIPASDTQQLAQEARNIYQEFLLSQALSPVNIDRQA
WLGEEVLAERPRDMFRAQQQLQIFNLMKFDSYARFVKSPLYRECLLAEAEGRPLREPGSSRLGSPDATRKK
PKLKPGKSLPLGVEELGQLPPVEGPGGRPLRKSFRRELGGTANAALRRESQGSLSNASSALDLGFLAFVSS
KSESHRKSLSGTEGESESRPGKYCCVYLPDGTASLALARPGLTIRDMLAGICEKRGSLSPDIKVVYLVGNE
QALVLDQDCTVLADQEVRLNRTFELELTALERVVRISAKPTKRLQEALQPILEKHGLSPLEVLHRPG
EKQPLDLGKLVSSVAAQRLVLDLTPGVKISKARDKSPCRSQGCPPTQDKATHPPASPSSLVKVPSSAT
GKRQTCDIEGLVELLNRVQSSGAHDQRGLLRKEDLVLPFLQLPAQGPSSEETPPQTKSAAQPIGGSLNS
TTDSAL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	61.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_006471
Locus ID:	10636
UniProt ID:	O43566
RefSeq Size:	2418
Cytogenetics:	5q35.3
RefSeq ORF:	1698
Summary:	<p>This gene encodes a member of the regulator of G-protein signaling family. This protein contains one RGS domain, two Raf-like Ras-binding domains (RBDs), and one GoLoco domain. The protein attenuates the signaling activity of G-proteins by binding, through its GoLoco domain, to specific types of activated, GTP-bound G alpha subunits. Acting as a GTPase activating protein (GAP), the protein increases the rate of conversion of the GTP to GDP. This hydrolysis allows the G alpha subunits to bind G beta/gamma subunit heterodimers, forming inactive G-protein heterotrimers, thereby terminating the signal. Alternate transcriptional splice variants of this gene have been observed but have not been thoroughly characterized. [provided by RefSeq, Jul 2008]</p>
Protein Families:	Druggable Genome

Product images:



Coomassie blue staining of purified RGS14 protein (Cat# TP304139). The protein was produced from HEK293T cells transfected with RGS14 cDNA clone (Cat# [RC204139]) using MegaTran 2.0 (Cat# [TT210002]).