

Product datasheet for TP304156M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Superoxide Dismutase 3 (SOD3) (NM 003102) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human superoxide dismutase 3, extracellular (SOD3), 100 μg

Species: Human
Expression Host: HEK293T

Expression cDNA >RC2
Clone or AA Red=
Sequence:

>RC204156 protein sequence Red=Cloning site Green=Tags(s)

MLALLCSCLLLAAGASDAWTGEDSAEPNSDSAEWIRDMYAKVTEIWQEVMQRRDDDGTLHAACQVQPSAT LDAAQPRVTGVVLFRQLAPRAKLDAFFALEGFPTEPNSSSRAIHVHQFGDLSQGCESTGPHYNPLAVPHP QHPGDFGNFAVRDGSLWRYRAGLAASLAGPHSIVGRAVVVHAGEDDLGRGGNQASVENGNAGRRLACCVV

GVCGPGLWERQAREHSERKKRRRESECKAA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 24.1 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling

conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 003093

Locus ID: 6649

UniProt ID: P08294, A0A140VJU8





RefSeq Size: 1546

Cytogenetics: 4p15.2
RefSeq ORF: 720
Synonyms: EC-SOD

Summary: This gene encodes a member of the superoxide dismutase (SOD) protein family. SODs are


antioxidant enzymes that catalyze the conversion of superoxide radicals into hydrogen peroxide

and oxygen, which may protect the brain, lungs, and other tissues from oxidative stress. Proteolytic processing of the encoded protein results in the formation of two distinct homotetramers that differ in their ability to interact with the extracellular matrix (ECM). Homotetramers consisting of the intact protein, or type C subunit, exhibit high affinity for heparin and are anchored to the ECM. Homotetramers consisting of a proteolytically cleaved form of the protein, or type A subunit, exhibit low affinity for heparin and do not interact with the ECM. A mutation in this gene may be associated with increased heart disease risk. [provided

by RefSeq, Oct 2015]

Protein Families: Druggable Genome, Secreted Protein

Product images:



Coomassie blue staining of purified SOD3 protein (Cat# [TP304156]). The protein was produced from HEK293T cells transfected with SOD3 cDNA clone (Cat# [RC204156]) using MegaTran 2.0 (Cat# [TT210002]).