

Product datasheet for TP304235L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DHPS (NM_001930) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human deoxyhypusine synthase (DHPS), transcript variant 1, 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC204235 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MEGSLEREAPAGALAAVLKHSSTLPPESTQVRGYDFNRGVNYRALLEAFGTTGFQATNFGRAVQQVNAMI EKKLEPLSQDEDQHADLTQSRRPLTSCTIFLGYTSNLISSGIRETIRYLVQHNMVDVLVTTAGGVEEDLI KCLAPTYLGEFSLRGKELRENGINRIGNLLVPNENYCKFEDWLMPILDQMVMEQNTEGVKWTPSKMIARL GKEINNPESVYYWAQKNHIPVFSPALTDGSLGDMIFFHSYKNPGLVLDIVEDLRLINTQAIFAKCTGMII LGGGVVKHHIANANLMRNGADYAVYINTAQEFDGSDSGARPDEAVSWGKIRVDAQPVKVYADASLVFPLL

VAETFAQKMDAFMHEKNED

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 40.8 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001921

Locus ID: 1725





UniProt ID: P49366, A0A024R7D0

RefSeq Size: 1361

Cytogenetics: 19p13.13 1107 RefSeq ORF:


Synonyms: DHS; DS; MIG13; NEDSSWI

Summary: This gene encodes a protein that is required for the formation of hypusine, a unique amino

acid formed by the posttranslational modification of only one protein, eukaryotic translation initiation factor 5A. The encoded protein catalyzes the first step in hypusine formation by transferring the butylamine moiety of spermidine to a specific lysine residue of the eukaryotic translation initiation factor 5A precursor, forming an intermediate deoxyhypusine residue. Alternatively spliced transcript variants encoding multiple isoforms have been observed for

this gene. [provided by RefSeq, May 2011]

Product images:



Coomassie blue staining of purified DHPS protein (Cat# [TP304235]). The protein was produced from HEK293T cells transfected with DHPS cDNA clone (Cat# [RC204235]) using MegaTran 2.0 (Cat# [TT210002]).