

Product datasheet for **TP304279M**

FURIN (NM_002569) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human furin (paired basic amino acid cleaving enzyme) (FURIN), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC204279 protein sequence Red =Cloning site Green =Tags(s)

MELRPWLLWVVAATGTLVLLAADAQGQKVFTNTWAVRIPGGPAVANSVARKHGFLNLGQIFGDYYHFWHR
GVTKRSLSPHRPRHSRLQREPQVQWLEQQVAKRRRTKRDVYQEPTDPKFPQQWYLSGVTQRDLNVKAAWAQ
GYTGHGIVWSILDDGIEKNHPDLAGNYDPGASFDVNDQDPDPQPRYTQMNDNRHGTRCAGEVAAVANNGV
CGVGVAYNARIGGVRMLDGEVTDAVEARSLGLNPNHIHYSASWGPEDDGKTVDGPALAEAEAFFRGVSQ
GRGGLGSIFVWASGNGGREHDCSCNDGYTNSIYTLSSSATQFGNVPWYSEACSTLATTYSSGNQNEKQ
IVTTDLRQKCTESHTGTSASAPLAAGIIALTLEANKNLTWRDMQHLLVQTSKPAHLNANDWATNGVGRKV
SHSYGYGLLDAGAMVALAQNWTTVAPQRKCIIDILTEPKDIGRLEVRKTVTACLGEPNHITRLEHAQAR
LTLSYNRRGDLAIHLVSPMGTRSTLLAARPHDYSADGFNDWAFMTTHSWDEDPGSEWVLEIENTSEANNY
GTLTKFTLVLYGTAPEGLPVPPESSGCKLTSSQACVCEEFGSLHQKSCVQHCPPGFAPQVLDTHYSTE
NDVETIRASVCAPCHASCATCQGPALTDCLSCPSHASLDPVEQTCSRQSQSSRESPPQQPPRLPPEVEA
GQRLRAGLLPSHLPEVVAGLSCAFIVLVFVTVFLVLQLRSGFSFRGVKVTMDRGLISYKGLPPEAWQEE
CPSDSEDEGRGERTAFIKDQSAL

TRTRPLEQLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	86.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.



[View online »](#)

Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_002560
Locus ID:	5045
UniProt ID:	P09958 , A0A024RC70
RefSeq Size:	4251
Cytogenetics:	15q26.1
RefSeq ORF:	2382
Synonyms:	FUR; PACE; PCSK3; SPC1
Summary:	<p>This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. It encodes a type 1 membrane bound protease that is expressed in many tissues, including neuroendocrine, liver, gut, and brain. The encoded protein undergoes an initial autocatalytic processing event in the ER and then sorts to the trans-Golgi network through endosomes where a second autocatalytic event takes place and the catalytic activity is acquired. Like other members of this convertase family, the product of this gene specifically cleaves substrates at single or paired basic residues. Some of its substrates include parathyroid hormone, transforming growth factor beta 1 precursor, proalbumin, pro-beta-secretase, membrane type-1 matrix metalloproteinase, beta subunit of pro-nerve growth factor and von Willebrand factor. It is thought to be one of the proteases responsible for the activation of HIV envelope glycoproteins gp160 and gp140, and may play a role in tumor progression. Unlike SARS-CoV and other coronaviruses, the spike protein of SARS-CoV-2 is thought to be uniquely cleaved by this protease. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2020]</p>
Protein Families:	Druggable Genome, Protease, Transmembrane

Product images:



Coomassie blue staining of purified FURIN protein (Cat# [TP304279]). The protein was produced from HEK293T cells transfected with FURIN cDNA clone (Cat# [RC204279]) using MegaTran 2.0 (Cat# [TT210002]).