

Product datasheet for **TP304333M**

UPF3A (NM_080687) Human Recombinant Protein


Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human UPF3 regulator of nonsense transcripts homolog A (yeast) (UPF3A), transcript variant 2, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC204333 protein sequence Red =Cloning site Green =Tags(s) MRSEKEGAGGLRAAVAARGPSGREKLSALEVQFHRDSQQQEAETPPTSSSGCGGGAGKPREEKRTALSKV VIRRLPPGLTKEQLEELRPLPAHDYFEFFAADLSLYPHLYSRAYINFRNPDDILLFRDRFDGYIFLDSK DPEYKKFLETYCVEEKTSANPETLLGEMEAKTRELIARRTTPLLEYIKNRKLEKQRIREEKREERRRRE LEKKRLREEEKRRRREEERCKKKETDKQKKIAEKEVRIKLLKKPEKGEEPTTEKPKERGEIDTGGGKQE SCAPGAVVKARPMEGSLEEPQETSHSGSDKEHRDVERSQQESEAQRYHVDDGRRHRAHHEPERLSRRS E DEQRWGKGPGQDRGKKGSQDSGAPGEAMERLGRAQRCDDSPAPRKERLANKDRPALQLYDPGARFRA REC GGNRRICKAEGSGTGPEKREEAE TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	50.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.


[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_542418</u>
Locus ID:	65110
UniProt ID:	<u>Q9H1J1</u>
RefSeq Size:	2303
Cytogenetics:	13q34
RefSeq ORF:	1329
Synonyms:	HUPF3A; RENT3A; UPF3
Summary:	This gene encodes a protein that is part of a post-splicing multiprotein complex involved in both mRNA nuclear export and mRNA surveillance. The encoded protein is one of two functional homologs to yeast Upf3p. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD to degrade mRNAs containing premature stop codons. This protein binds to the mRNA and remains bound after nuclear export, acting as a nucleocytoplasmic shuttling protein. It forms with Y14 a complex that binds specifically 20 nt upstream of exon-exon junctions. This gene is located on the long arm of chromosome 13. Several splice variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2017]

Product images:



Coomassie blue staining of purified UPF3A protein (Cat# [TP304333]). The protein was produced from HEK293T cells transfected with UPF3A cDNA clone (Cat# [RC204333]) using MegaTran 2.0 (Cat# [TT210002]).