

Product datasheet for **TP304369M**

BAX (NM_138761) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens BCL2-associated X protein (BAX), transcript variant alpha, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC204369 representing NM_138761 Red =Cloning site Green =Tags(s)
	MDGSGEQPRGGGPTSSEQIMKTGALLLQGFQDRAGRMGGAEPELALDPVPQDASTKKLSECLKRIGDEL DSNMELQRMIAAVDTDSPREVFFRVAADMFSNGNFWGRVVALFYFASKLVLKALCTKVPELIRTIMGWT LDFLRERLLGWIQDQGGWDGLLSYFGTPTWQTVTIFVAGVLTASLTIWKKMG
	TR TRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	21 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_620116</u>
Locus ID:	581
UniProt ID:	<u>Q07812</u>



[View online »](#)

RefSeq Size: 888

Cytogenetics: 19q13.33

RefSeq ORF: 576


Synonyms: BCL2L4

Summary: The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein forms a heterodimer with BCL2, and functions as an apoptotic activator. The association and the ratio of BAX to BCL2 also determines survival or death of a cell following an apoptotic stimulus. This protein is reported to interact with, and increase the opening of, the mitochondrial voltage-dependent anion channel (VDAC), which leads to the loss in membrane potential and the release of cytochrome c. The expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis. Multiple alternatively spliced transcript variants, which encode different isoforms, have been reported for this gene. [provided by RefSeq, Dec 2019]

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Amyotrophic lateral sclerosis (ALS), Apoptosis, Colorectal cancer, Huntington's disease, Neurotrophin signaling pathway, p53 signaling pathway, Pathways in cancer, Prion diseases

Product images:



Coomassie blue staining of purified BAX protein (Cat# [TP304369]). The protein was produced from HEK293T cells transfected with BAX cDNA clone (Cat# [RC204369]) using MegaTran 2.0 (Cat# [TT210002]).