

Product datasheet for **TP304442M**

CDK2AP1 (NM_004642) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human cyclin-dependent kinase 2 associated protein 1 (CDK2AP1), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC204442 protein sequence Red =Cloning site Green =Tags(s)
	MSYKPNLAAHMPAAALNAAGSVHSPSTSMATSSQYRQLLSYDGPPSLGYTQGTGNSQVPQSKYAELLAI EELGKEIRPTYAGSKSAMERLKRGIHARGLVRECLAETERNARS
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	12.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_004633
Locus ID:	8099
UniProt ID:	O14519
RefSeq Size:	1655



[View online »](#)


Cytogenetics: 12q24.31

RefSeq ORF: 345

Synonyms: doc-1; DOC1; DORC1; p12DOC-1; ST19

Summary: The protein encoded by this gene is a cyclin-dependent kinase 2 (CDK2) -associated protein which is thought to negatively regulate CDK2 activity by sequestering monomeric CDK2, and targeting CDK2 for proteolysis. This protein was found to also interact with DNA polymerase alpha/primase and mediate the phosphorylation of the large p180 subunit, which suggests a regulatory role in DNA replication during the S-phase of the cell cycle. This protein also forms a core subunit of the nucleosome remodeling and histone deacetylation (NURD) complex that epigenetically regulates embryonic stem cell differentiation. This gene thus plays a role in both cell-cycle and epigenetic regulation. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2012]

Product images:



Coomassie blue staining of purified CDK2AP1 protein (Cat# [TP304442]). The protein was produced from HEK293T cells transfected with CDK2AP1 cDNA clone (Cat# [RC204442]) using MegaTran 2.0 (Cat# [TT210002]).