

## **Product datasheet for TP304552**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### HEXIM2 (NM\_144608) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human hexamthylene bis-acetamide inducible 2 (HEXIM2), 20 μg

Species: Human Expression Host: HEK293T

**Expression cDNA Clone** >RC204552 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MMATPNQTACNAESPVALEEAKTSGAPGSPQTPPERHDSGGSLPLTPRMESHSEDEDLAGAVGGLGWNSR SPRTQSPGGCSAEAVLARKKHRRPSKRKRHWRPYLELSWAEKQQRDERQSQRASRVREEMFAKGQPVAP YNTTQFLMNDRDPEEPNLDVPHGISHPGSSGESEAGDSDGRGRAHGEFQRKDFSETYERFHTESLQGRSK QELVRDYLELEKRLSQAEEETRRLQQLQACTGQQSCRQVEELAAEVQRLRTENQRLRQENQMWNREGCRC

DEEPGT

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK
Predicted MW: 32.2 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling

conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 653209 **Locus ID:** 124790



Synonyms:

#### HEXIM2 (NM\_144608) Human Recombinant Protein - TP304552

**UniProt ID:** Q96MH2

1330 RefSeq Size:

Cytogenetics: 17q21.31

RefSeq ORF: 858 L3


**Summary:** This gene encodes a member of the HEXIM family of proteins. This protein is a component of

> the 7SK small nuclear ribonucleoprotein. This protein has been found to negatively regulate the kinase activity of the cyclin-dependent kinase P-TEFb, which phosphorylates multiple target proteins to promote transcriptional elongation. This gene is located approximately 7 kb downstream from related family member HEXIM1 on chromosome 17. Alternative splicing

results in multiple transcript variants. [provided by RefSeq, Jan 2015]

**Protein Families: Transcription Factors** 

# **Product images:**



Coomassie blue staining of purified HEXIM2 protein (Cat# TP304552). The protein was produced from HEK293T cells transfected with HEXIM2 cDNA clone (Cat# [RC204552]) using MegaTran 2.0 (Cat# [TT210002]).