

Product datasheet for **TP304636**

CCN5 (NM_003881) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human WNT1 inducible signaling pathway protein 2 (WISP2), 20 µg

Species: Human

Expression Host: HEK293T

Expression cDNA >RC204636 protein sequence

Clone or AA Sequence: **Red**=Cloning site **Green**=Tags(s)

MRGTPKTHLLAFSLLCLLSKVRTQLCPTPCTCPWPPRCPLGVPLVLDGCGCCRVCARRLGEPDQLHVC
DASQGLVCQPGAGPGGRGALCLLAEDDSSCEVNGRLYREGETFQPHCSIRCRCEGGFTCVPLCSEDVRL
PSWDCPHRRVEVLGKCCPEWVCGQGGGLGTQPLPAQGPQFSGLVSSLPPGVPCPEWSTAWGPCSTTCGL
GMATRVSNQNRFCRLETQRRLLCLSRPCPPSRGRSPQNSAF

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 24.2 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: [NP_003872](#)

Locus ID: 8839

UniProt ID: [O76076](#)



[View online >](#)

RefSeq Size: 1433

Cytogenetics: 20q13.12


RefSeq ORF: 750

Synonyms: CT58; CTGF-L; WISP2

Summary: This gene encodes a member of the WNT1 inducible signaling pathway (WISP) protein subfamily, which belongs to the connective tissue growth factor (CTGF) family. WNT1 is a member of a family of cysteine-rich, glycosylated signaling proteins that mediate diverse developmental processes. The CTGF family members are characterized by four conserved cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C module, thrombospondin domain and C-terminal cystine knot-like (CT) domain. The encoded protein lacks the CT domain which is implicated in dimerization and heparin binding. It is 72% identical to the mouse protein at the amino acid level. This gene may be downstream in the WNT1 signaling pathway that is relevant to malignant transformation. Its expression in colon tumors is reduced while the other two WISP members are overexpressed in colon tumors. It is expressed at high levels in bone tissue, and may play an important role in modulating bone turnover. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein

Product images:



Coomassie blue staining of purified CCN5 protein (Cat# TP304636). The protein was produced from HEK293T cells transfected with CCN5 cDNA clone (Cat# [RC204636]) using MegaTran 2.0 (Cat# [TT210002]).