

## **Product datasheet for TP304709**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Hemoglobin subunit gamma 2 (HBG2) (NM 000184) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human hemoglobin, gamma G (HBG2), 20 μg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC204709 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MGHFTEEDKATITSLWGKVNVEDAGGETLGRLLVVYPWTQRFFDSFGNLSSASAIMGNPKVKAHGKKVLT SLGDAIKHLDDLKGTFAQLSELHCDKLHVDPENFKLLGNVLVTVLAIHFGKEFTPEVQASWQKMVTAVAS

**ALSSRYH** 

**TRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 

Tag: C-Myc/DDK
Predicted MW: 15.9 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 000175

 Locus ID:
 3048

 UniProt ID:
 P69892

 RefSeq Size:
 583



Cytogenetics: 11p15.4

RefSeq ORF: 441

Synonyms: HBG-T1; TNCY


Summary: The gamma globin genes (HBG1 and HBG2) are normally expressed in the fetal liver, spleen

and bone marrow. Two gamma chains together with two alpha chains constitute fetal hemoglobin (HbF) which is normally replaced by adult hemoglobin (HbA) at birth. In some

beta-thalassemias and related conditions, gamma chain production continues into adulthood. The two types of gamma chains differ at residue 136 where glycine is found in the G-gamma product (HBG2) and alanine is found in the A-gamma product (HBG1). The former is predominant at birth. The order of the genes in the beta-globin cluster is: 5'- epsilon --

gamma-G -- gamma-A -- delta -- beta--3'. [provided by RefSeq, Jul 2008]

## **Product images:**



Coomassie blue staining of purified HBG2 protein (Cat# TP304709). The protein was produced from HEK293T cells transfected with HBG2 cDNA clone (Cat# [RC204709]) using MegaTran 2.0 (Cat# [TT210002]).