

## Product datasheet for **TP304788**

### EIF3E (NM\_001568) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human eukaryotic translation initiation factor 3, subunit E (EIF3E), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC204788 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MAEYDLTTRIAHFLDRHLVFPLEFLSVKEIYNEKELLQGKLDLLSDTNMVFAMDVYKNLYSDDIPHAL  
REKRTTVVAQLKQLQAETEPVIMFEDPETTRQMQRDGRMLFDYLADKHGFRQEYLDLTRYAKFQYE  
CGNYSGAAEYLYFFRVLVPATDRNALSSLWGKLASEILMQNWDVAVMEDLTRLKETIDNNSVSSPLQSLQQ  
RTWLIHWSLFFVFNHPKGRDNIIDLFLYQPQYLNAIQTMCPHILRYLTTAVITNKDVRKRRQVLKDLVKV  
IQQESYTYKDPITEFVECLYVNFDFDGAQKKLRECESVLVNDFFLVACLEDFIENARLFIFETFCRIHQ  
ISINMLADKLNMTPEEAERWIVNLIRNARLDAKIDSKLGHVVMGNNAVSPYQQVIEKTKSLSFRSQMLAM  
NIEKLNQNSRSEAPNWATQDSGFY

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**

Tag:	C-Myc/DDK
Predicted MW:	52 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.




[View online »](#)

RefSeq:	<a href="#">NP_001559</a>
Locus ID:	3646
UniProt ID:	<a href="#">P60228</a>
RefSeq Size:	1516
Cytogenetics:	8q23.1
RefSeq ORF:	1335
Synonyms:	eIF3-p46; EIF3-P48; EIF3S6; INT6

**Summary:** Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:17581632, PubMed:25849773, PubMed:27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA<sup>i</sup> and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:17581632). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:25849773). Required for nonsense-mediated mRNA decay (NMD); may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway (PubMed:17468741). May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins (PubMed:17310990, PubMed:17324924). [UniProtKB/Swiss-Prot Function]

### Product images:



Coomassie blue staining of purified EIF3E protein (Cat# TP304788). The protein was produced from HEK293T cells transfected with EIF3E cDNA clone (Cat# [RC204788]) using MegaTran 2.0 (Cat# [TT210002]).