

Product datasheet for **TP304869L**

IRAK (IRAK1) (NM_001025243) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human interleukin-1 receptor-associated kinase 1 (IRAK1), transcript variant 3, 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC204869 protein sequence Red =Cloning site Green =Tags(s)

MAGGPGPGEPAAPGAQHFLYEVPPWVMCRFYKVMDALEPADWCQFAALIVRDQTELRRLCERSGQRTASVL
WPWINRNARVADLVHILHLQLLRARDIITAWHPPAPLPSPGTTAPRPSSIPAPAEAEAWSPRKLPSAS
TFLSPAFIGSQTHSGPELGLVPSASLWPPPPSPAPSSTKPGPESSVSLQGARPSPCWPLCEISRGT
NFSEELKIGEGGFGCVYRAVMRNTVYAVKRLKENADLEWTAVKQSFLTEVEQLSRFRHPNIVDFAGYCAQ
NGFYCLVYGFLPNGSLEDRLHCQTQACPLSWPQRDILLGTARAIQFLHQDSPSLIHGDIKSSNVLLDE
RLTPKLGDFGLARFSRFAGSSPSQSSMVARTQTVRGTLAYLPEEYIKTGRLAVDTDTFSGVWVLETLAG
QRAVKTHGARTKYLVERLEKLQAVVAGVPGHLEAASCIPPSPQENSYSSTGRAHSGAAPWQPLAAPSG
ASAQAAEQLQRGNQPVESESLGGLSAAALRSWHLTPSCPLDPAPLREAGCPQGD TAGESSWGS GPGSRP
TAVEGLALGSSASSSSEPPQIIINPARQKMVQKLALYEDGALDSLQLLSSSSLPGLGLEQDRQGPEESDE
FQS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	67.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_001020414
Locus ID:	3654
UniProt ID:	P51617
RefSeq Size:	3352
Cytogenetics:	Xq28
RefSeq ORF:	1899
Synonyms:	IRAK; pelle
Summary:	This gene encodes the interleukin-1 receptor-associated kinase 1, one of two putative serine/threonine kinases that become associated with the interleukin-1 receptor (IL1R) upon stimulation. This gene is partially responsible for IL1-induced upregulation of the transcription factor NF-kappa B. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
Protein Families:	Druggable Genome, Protein Kinase, Transcription Factors
Protein Pathways:	Apoptosis, Neurotrophin signaling pathway, Toll-like receptor signaling pathway

Product images:



Coomassie blue staining of purified IRAK1 protein (Cat# [TP304869]). The protein was produced from HEK293T cells transfected with IRAK1 cDNA clone (Cat# [RC204869]) using MegaTran 2.0 (Cat# [TT210002]).