

## **Product datasheet for TP304872L**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## XPA (NM\_000380) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human xeroderma pigmentosum, complementation group A

(XPA), transcript variant 1, full length, with C-terminal MYC/DDK tag, expressed in HEK293 cells,

1 mg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC204872 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MAAADGALPEAAALEQPAELPASVRASIERKRQRALMLRQARLAARPYSATAAAATGGMANVKAAPKIID TGGGFILEEEEEEQKIGKVVHQPGPVMEFDYVICEECGKEFMDSYLMNHFDLPTCDNCRDADDKHKLIT KTEAKQEYLLKDCDLEKREPPLKFIVKKNPHHSQWGDMKLYLKLQIVKRSLEVWGSQEALEEAKEVRQEN REKMKQKKFDKKVKELRRAVRSSVWKRETIVHQHEYGPEENLEDDMYRKTCTMCGHELTYEKM

**TRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 

Tag: C-Myc/DDK

Predicted MW: 31.2 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

**Storage:** Store at -80°C.

**Stability:** Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 000371

**Locus ID:** 7507





UniProt ID: P23025

RefSeq Size: 1491

**Cytogenetics:** 9q22.33

RefSeq ORF: 819

Synonyms: XP1; XPAC

Summary: This gene encodes a zinc finger protein plays a central role in nucleotide excision repair (NER),


a specialized type of DNA repair. NER is responsible for repair of UV radiation-induced photoproducts and DNA adducts induced by chemical carcinogens and chemotherapeutic drugs. The encoded protein interacts with DNA and several NER proteins, acting as a scaffold to assemble the NER incision complex at sites of DNA damage. Mutations in this gene cause Xeroderma pigmentosum complementation group A (XP-A), an autosomal recessive skin disorder featuring hypersensitivity to sunlight and increased risk for skin cancer. [provided by

RefSeq, Aug 2017]

**Protein Families:** Druggable Genome

**Protein Pathways:** Nucleotide excision repair

## **Product images:**



Coomassie blue staining of purified XPA protein (Cat# [TP304872]). The protein was produced from HEK293T cells transfected with XPA cDNA clone (Cat# [RC204872]) using MegaTran 2.0 (Cat# [TT210002]).