

Product datasheet for TP304932

Tbp7 (PSMC4) (NM_006503) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human proteasome (prosome, macropain) 26S subunit, ATPase, 4 (PSMC4), transcript variant 1, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC204932 protein sequence Red=Cloning site Green=Tags(s)

MEEIGILVEKAQDEIPALSVSRPQTGLSFLGPEPEDLEDLYSRYKKLQQEFLFLEVQEEYIKDEQKNLKK
EFLHAQEEVKRIQSIPLVIGQFLEAVDQNTAIVGTTGNSYVYRILSTIDRELLKPNASVALHKHSNALV
DVLPPHADSSIMMLTSDQKPDVMYADIGGMDIQKQEVREAVELPLTHFELYKQIGIDPPRGVLMYGPPGC
GKTMLAKAVAHHTTAAFIRVVGSEFVQKYLGEPRMVRDVFLAKENAPAIIFIDEIDAIATKRFDAQTG
ADREVQRILLELLNQMDGFDQNVNVKIMATNRADTLDPALLRPGRLDRKIEFPLPDRRQKRLIFSTITS
KMNLSEEVDLEDYVARPDKISGADINSICQESGMLAVRENRYIVLAKDFEKAYKTVIKKDEQEHEFYK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	47.2 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_006494



[View online »](#)

Locus ID: 5704

UniProt ID: [P43686](#), [A8K2M0](#)

RefSeq Size: 1914

Cytogenetics: 19q13.2

RefSeq ORF: 1254

Synonyms: MIP224; RPT3; S6; TBP-7; TBP7

Summary: The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. This gene encodes a member of the triple-A family of ATPases that is a component of the 19S regulatory subunit and plays a role in 26S proteasome assembly. The encoded protein interacts with gankyrin, a liver oncoprotein, and may also play a role in Parkinson's disease through interactions with synphilin-1. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jul 2012]

Protein Families: Druggable Genome

Protein Pathways: Proteasome

Product images:



Coomassie blue staining of purified PSMC4 protein (Cat# TP304932). The protein was produced from HEK293T cells transfected with PSMC4 cDNA clone (Cat# [RC204932]) using MegaTran 2.0 (Cat# [TT210002]).