

Product datasheet for TP305066L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Viperin (RSAD2) (NM_080657) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human radical S-adenosyl methionine domain containing 2 (RSAD2),

1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

>RC205066 protein sequence Red=Cloning site Green=Tags(s)

MWVLTPAAFAGKLLSVFRQPLSSLWRSLVPLFCWLRATFWLRATKRRKQQLVLRGPDETKEEEEDPPLPT TPTSVNYHFTRQCNYKCGFCFHTAKTSFVLPLEEAKRGLLLLKEAGMEKINFSGGEPFLQDRGEYLGKLV RFCKVELRLPSVSIVSNGSLIRERWFQNYGEYLDILAISCDSFDEEVNVLIGRGQGKKNHVENLQKLRRW CRDYRVAFKINSVINRFNVEEDMTEQIKALNPVRWKVFQCLLIEGENCGEDALREAERFVIGDEEFERFL ERHKEVSCLVPESNQKMKDSYLILDEYMRFLNCRKGRKDPSKSILDVGVEEAIKFSGFDEKMFLKRGGKY

IWSKADLKLDW

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 42 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 542388



Locus ID: 91543

UniProt ID: Q8WXG1

RefSeq Size: 3512 Cytogenetics: 2p25.2 RefSeq ORF: 1083


Synonyms: cig5; cig33; vig1

Summary: The protein encoded by this gene is an interferon-inducible antiviral protein that belongs to

the S-adenosyl-L-methionine (SAM) superfamily of enzymes. The protein plays a role in cellular antiviral response and innate immune signaling. Antiviral effects result from inhibition of viral RNA replication, interference in the secretory pathway, binding to viral proteins and dysregulation of cellular lipid metabolism. The protein has been found to inhibit both DNA and RNA viruses, including influenza virus, human immunodeficiency virus (HIV-1) and Zika

virus. [provided by RefSeq, Sep 2020]

Product images:



Coomassie blue staining of purified RSAD2 protein (Cat# [TP305066]). The protein was produced from HEK293T cells transfected with RSAD2 cDNA clone (Cat# [RC205066]) using MegaTran 2.0 (Cat# [TT210002]).