

Product datasheet for **TP305212M**

SUCLA2 (NM_003850) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human succinate-CoA ligase, ADP-forming, beta subunit (SUCLA2), nuclear gene encoding mitochondrial protein, 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC205212 protein sequence Red =Cloning site Green =Tags(s) MAASMFYGRLLVAVATLRNHRPRTAQRAAAQVLGSSGLFNNHGLQVQQQQQRNLSLHEYMSMELLQEAGVS VPKGYVAKSPDEAYAIKKLGSKDVWIKAQVLAGGRGKGTTFESGLKGGVKIVFSPEEAKAVSSQMIGKKL FTKQTGEKGRICNQVLVCERKYPREYYFAITMERSFQGPVLIGSSHGGVNIEDVAAESPEAIKEPIDI EEGIKKEQALQLAQKMGFPPNIVESAAENMVKLYSLFLKYDATMIEINPMVEDSDGAVLCMDAKINFDSN SAYRQKKIFDLQDWTQEDERDKDAAKANLNYIGLDGNIGCLVNGAGLAMATMDIIKLHGGTPANFLDVGG GATVHQVTEAFKLITSDKKVLAILVNIFFGGIMRCDVIAQGIVMAVKDLEIKIPVVVRLQGTRVDDAKALI ADSGLKILACDDLDEAARMVVKLSEIVTLAKQAHVDVKFQLPI TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	44.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_003841](#)

Locus ID: 8803

UniProt ID: [Q9P2R7](#), [E5KS60](#), [Q9Y4T0](#)

RefSeq Size: 2182

Cytogenetics: 13q14.2

RefSeq ORF: 1389

Synonyms: A-BETA; A-SCS; LINC00444; MTDPS5; SCS-betaA

Summary: Succinyl-CoA synthetase (SCS) is a mitochondrial matrix enzyme that acts as a heterodimer, being composed of an invariant alpha subunit and a substrate-specific beta subunit. The protein encoded by this gene is an ATP-specific SCS beta subunit that dimerizes with the SCS alpha subunit to form SCS-A, an essential component of the tricarboxylic acid cycle. SCS-A hydrolyzes ATP to convert succinate to succinyl-CoA. Defects in this gene are a cause of myopathic mitochondrial DNA depletion syndrome. A pseudogene of this gene has been found on chromosome 6. [provided by RefSeq, Jul 2008]

Protein Pathways: Citrate cycle (TCA cycle), Metabolic pathways, Propanoate metabolism

Product images:



Coomassie blue staining of purified SUCLA2 protein (Cat# [TP305212]). The protein was produced from HEK293T cells transfected with SUCLA2 cDNA clone (Cat# [RC205212]) using MegaTran 2.0 (Cat# [TT210002]).