

Product datasheet for **TP305238M**

TBK1 (NM_013254) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human TANK-binding kinase 1 (TBK1), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC205238 protein sequence Red =Cloning site Green =Tags(s)

MQSTSNHLWLLSDILGQGATANVFRGRHKKTGDLFAIKVFNNISFLRPVDVQMREFEVLKKNLHNKIVKL
FAIEEETTRHKVLIIMEFCPCGSLYTVLEEPSNAYGLPESEFLIVLRDVGGMNHLRENGIVHRDIKPGN
IMRVIGEDGQSUYKLTDFGAARELEDDEQFVSLYGTEEYLHPDMYERAVLRKDHQKKYGATVDLWSIGVT
FYHAATGSLPFRPFEGPRRNKEVMYKIITGKPSGAISGVQKAENGPIDWSGDMPVSCSLRGLQVLLTPV
LANILEADQEKCWGFDDQFAETSDILHRMVIHVFSLQQMTAHKIYIHSYNTATIFHELVIKQTKIISNQ
ELIYEGRRLLVLEPGRLAQHFPKTTEENPIFVVSREPLDTIGLIYEKISLPKVHPRYDLGDASMAKAITG
VVCYACRIASTLLLYQELMRKGIWLIELIKDDYNETVHKKTEVITLDFCIRNIEKTVKVEKLMKINL
EAAELGEISDIHTKLLRLSSSQGTIETSLQDIDSRLSPGGSLADAWAHQEGTHPKDRNVEKLQVLLNCMT
EIYYQFKKDAQERRLAYNEEQIHKFDKQKLYHATKAMTHFTDECVKKYEAFLNKSEEWIRKMLHLRKQL
LSLTNQCFDIEEEVSKYQEYTNELQETLPQKMFTASSGIKHTMTPIYSSNTLVEMTLGMKKLKEEMEGV
VKELAENNHILERFGSLTMDGGLRNVDCI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Tag:	C-Myc/DDK
Predicted MW:	83.5 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_037386
Locus ID:	29110
UniProt ID:	Q9UHD2
RefSeq Size:	3098
Cytogenetics:	12q14.2
RefSeq ORF:	2187
Synonyms:	FTDALS4; IIAE8; NAK; T2K
Summary:	The NF-kappa-B (NFKB) complex of proteins is inhibited by I-kappa-B (IKB) proteins, which inactivate NFKB by trapping it in the cytoplasm. Phosphorylation of serine residues on the IKB proteins by IKB kinases marks them for destruction via the ubiquitination pathway, thereby allowing activation and nuclear translocation of the NFKB complex. The protein encoded by this gene is similar to IKB kinases and can mediate NFKB activation in response to certain growth factors. [provided by RefSeq, Oct 2010]
Protein Families:	Druggable Genome, Protein Kinase
Protein Pathways:	Cytosolic DNA-sensing pathway, RIG-I-like receptor signaling pathway, Toll-like receptor signaling pathway

Product images:



Coomassie blue staining of purified TBK1 protein (Cat# [TP305238]). The protein was produced from HEK293T cells transfected with TBK1 cDNA clone (Cat# [RC205238]) using MegaTran 2.0 (Cat# [TT210002]).