

## **Product datasheet for TP305403**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## DDI1 (NM 001001711) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human DDI1, DNA-damage inducible 1, homolog 1 (S. cerevisiae)

(DDI1), 20 µg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC205403 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MLITVYCVRRDLSEVTFSLQVSPDFELRNFKVLCEAESRVPVEEIQIIHMERLLIEDHCSLGSYGLKDGD IVVLLQKDNVGPRAPGRAPNQPRVDFSGIAVPGTSSSRPQHPGQQQQRTPAAQRSQGLASGEKVAGLQ

GL

GSPALIRSMLLSNPHDLSLLKERNPPLAEALLSGSLETFSQVLMEQQREKALREQERLRLYTADPLDREA QAKIEEEIRQQNIEENMNIAIEEAPESFGQVTMLYINCKVNGHPLKAFVDSGAQMTIMSQACAERCNIMR LVDRRWAGVAKGVGTQRIIGRVHLAQIQIEGDFLQCSFSILEDQPMDMLLGLDMLRRHQCSIDLKKNVLV

IGTTGTQTYFLPEGELPLCSRMVSGQDESSDKEITHSVMDSGRKEH

**TRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 

Tag: C-Myc/DDK
Predicted MW: 43.9 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.





**RefSeq:** NP 001001711

 Locus ID:
 414301

 UniProt ID:
 Q8WTU0

 RefSeq Size:
 2615

 Cytogenetics:
 11q22.3


RefSeq ORF: 1188

Summary: Probable aspartic protease (Probable). Seems to act as a proteasomal shuttle which links the

proteasome and replication fork proteins like RTF2 (Probable). Required, with DDI2, for cellular survival following replication stress. Together or redudantly with DDI2, removes RTF2 from stalled forks to allow cell cycle progression after replication stress and maintains

genome integrity (PubMed:29290612).[UniProtKB/Swiss-Prot Function]

## **Product images:**



Coomassie blue staining of purified DDI1 protein (Cat# TP305403). The protein was produced from HEK293T cells transfected with DDI1 cDNA clone (Cat# [RC205403]) using MegaTran 2.0 (Cat# [TT210002]).