

## Product datasheet for **TP305403L**

### DDI1 (NM\_001001711) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human DDI1, DNA-damage inducible 1, homolog 1 ( <i>S. cerevisiae</i> ) (DDI1), 1 mg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC205403 protein sequence <span style="color: red;">Red</span> =Cloning site <span style="color: green;">Green</span> =Tags(s)  MLITVYCVRRDLSEVTFSLQVSPDFELRNFKVLCEASRPVVEEQIIHMERLLIEDHCSLSYGLKDGD IVVLLQKDNVGPAPGRAPNQPRVDFSGIAVPGTSSSRPQHPGQQQQRTPAQRSQGLASGEKVAGLQ GL GSPALIRSMLLSNPHDLSLLKERNPPLAEALLSGSLETFSQVLMEQQREKALREQERLRLYTADPLDREA QAKIEEEIRQQNIEENMNIAIEEAPESFGQVTMLYINCKVNGHPLKAFVDSGAQMTIMSQACAERCNIMR LVDRRWAGVAKGVGTQRIIGRVHLAQIQIEGDFLQCSFSILEDQPMMDMLLGLDMLRRHQCSIDLKKNVLV IGTGTGTQTYFLPEGELPLCSRMSVSGQDESSDKEITHSVMDSGRKEH  <span style="color: red;">TR</span> <span style="color: green;">TRPLEQKLISEEDLAANDILDYKDDDDKV</span>
Tag:	C-Myc/DDK
Predicted MW:	43.9 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.


[View online »](#)

RefSeq: NP\_001001711

Locus ID: 414301

UniProt ID: Q8WTU0


RefSeq Size: 2615

Cytogenetics: 11q22.3

RefSeq ORF: 1188

**Summary:** Probable aspartic protease (Probable). Seems to act as a proteasomal shuttle which links the proteasome and replication fork proteins like RTF2 (Probable). Required, with DDI2, for cellular survival following replication stress. Together or redudantly with DDI2, removes RTF2 from stalled forks to allow cell cycle progression after replication stress and maintains genome integrity (PubMed:29290612).[UniProtKB/Swiss-Prot Function]

## Product images:



Coomassie blue staining of purified DDI1 protein (Cat# [TP305403]). The protein was produced from HEK293T cells transfected with DDI1 cDNA clone (Cat# [RC205403]) using MegaTran 2.0 (Cat# [TT210002]).