

## **Product datasheet for TP305453**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HORMAD2 (NM\_152510) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human HORMA domain containing 2 (HORMAD2), 20 μg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC205453 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MATAQLSHCITIHKASKETVFPSQITNEHESLKMVKKLFATSISCITYLRGLFPESSYGERHLDDLSLKI LREDKKCPGSLHIIRWIQGCFDALEKRYLRMAVLTLYTDPMGSEKVTEMYQFKFKYTKEGATMDFDSHSS STSFESGTNNEDIKKASVLLIRKLYILMQDLEPLPNNVVLTMKLHYYNAVTPHDYQPLGFKEGVNSHFLL FDKEPINVQVGFVSTGFHSMKVKVMTEATKVIDLENNLFRENSTTEIAHQGLDCDEEEECNDHIQRMNFV

CSQQSSECSRKKRKVSEPVKVFIPNRK

**TRTRPL**EQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 35.1 kDa

Concentration: >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 689723 **Locus ID:** 150280






UniProt ID: Q8N7B1 1909 RefSeq Size: Cytogenetics: 22q12.2 RefSeq ORF: 921 Synonyms: CT46.2

**Summary:** Essential for synapsis surveillance during meiotic prophase via the recruitment of ATR activity.

Plays a key role in the male mid-pachytene checkpoint and the female meiotic prophase checkpoint: required for efficient build-up of ATR activity on unsynapsed chromosome regions, a process believed to form the basis of meiotic silencing of unsynapsed chromatin (MSUC) and meiotic prophase quality control in both sexes. Required for the DNA doublestrand break-independent, BRCA1-dependent activation of ATR on the sex chromosomes that is essential for normal sex body formation (By similarity).[UniProtKB/Swiss-Prot Function]

## **Product images:**



Coomassie blue staining of purified HORMAD2 protein (Cat# TP305453). The protein was produced from HEK293T cells transfected with HORMAD2 cDNA clone (Cat# [RC205453]) using MegaTran 2.0 (Cat# [TT210002]).