

Product datasheet for TP305607M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SAP18 (NM_005870) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human Sin3A-associated protein, 18kDa (SAP18), 100 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC205607 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MAVESRVTQEEIKKEPEKPIDREKTCPLLLRVFTTNNGRHHRMDEFSRGNVPSSELQIYTWMDATLKELT SLVKEVYPEARKKGTHFNFAIVFTDVKRPGYRVKEIGSTMSGRKGTDDSMTLQSQKFQIGDYLDIAITPP

NRAPPTSGRMRPY

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 17.4 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 005861

 Locus ID:
 10284

 UniProt ID:
 000422

 RefSeq Size:
 2318





Cytogenetics: 13q12.11

RefSeq ORF: 459

Synonyms: 2HOR0202; SAP18P


Summary: Histone acetylation plays a key role in the regulation of eukaryotic gene expression. Histone

acetylation and deacetylation are catalyzed by multisubunit complexes. The protein encoded by this gene is a component of the histone deacetylase complex, which includes SIN3, SAP30, HDAC1, HDAC2, RbAp46, RbAp48, and other polypeptides. This protein directly interacts with SIN3 and enhances SIN3-mediated transcriptional repression when tethered to the promoter.

A pseudogene has been identified on chromosome 2. [provided by RefSeq, Dec 2008]

Protein Families: Druggable Genome, Transcription Factors

Product images:



Coomassie blue staining of purified SAP18 protein (Cat# [TP305607]). The protein was produced from HEK293T cells transfected with SAP18 cDNA clone (Cat# [RC205607]) using MegaTran 2.0 (Cat# [TT210002]).