

## **Product datasheet for TP305615**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## BCAS2 (NM\_005872) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human breast carcinoma amplified sequence 2 (BCAS2), 20 μg

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** >RC205615 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MAGTGLVAGEVVVDALPYFDQGYEAPGVREAAAALVEEETRRYRPTKNYLSYLTAPDYSAFETDIMRNEF ERLAARQPIELLSMKRYELPAPSSGQKNDITAWQECVNNSMAQLEHQAVRIENLELMSQHGCNAWKVYNE NLVHMIEHAQKELQKLRKHIQDLNWQRKNMQLTAGSKLREMESNWVSLVSKNYEIERTIVQLENEIYQIK

QQHGEANKENIRQDF

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK

**Predicted MW:** 26 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 005863

**Locus ID:** 10286

**UniProt ID:** 075934, B2R7W3





RefSeq Size: 1310

Cytogenetics: 1p13.2 RefSeq ORF: 675

Synonyms: DAM1; Snt309; SPF27

**Summary:** Required for pre-mRNA splicing as component of the activated spliceosome


(PubMed:28502770, PubMed:28076346, PubMed:29360106, PubMed:29301961,

PubMed:30705154). Component of the PRP19-CDC5L complex that forms an integral part of the spliceosome and is required for activating pre-mRNA splicing. May have a scaffolding role in the spliceosome assembly as it contacts all other components of the core complex. The PRP19-CDC5L complex may also play a role in the response to DNA damage (DDR).

[UniProtKB/Swiss-Prot Function]

Spliceosome **Protein Pathways:** 

## **Product images:**



Coomassie blue staining of purified BCAS2 protein (Cat# TP305615). The protein was produced from HEK293T cells transfected with BCAS2 cDNA clone (Cat# [RC205615]) using MegaTran 2.0 (Cat# [TT210002]).