

Product datasheet for TP305676L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Troponin I fast skeletal muscle (TNNI2) (NM_003282) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human troponin I type 2 (skeletal, fast) (TNNI2), transcript variant 1, 1

mg

Species: Human Expression Host: HEK293T

Expression cDNA Clone >RC205676 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

LKQVKKEDTEKERDLRDVGDWRKNIEEKSGMEGRKKMFESES

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK
Predicted MW: 21.2 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 003273

Locus ID: 7136

UniProt ID: P48788





RefSeq Size: 738

Cytogenetics: 11p15.5 RefSeq ORF: 546


Synonyms: AMCD2B; DA2B; DA2B1; FSSV; fsTnl

Summary: This gene encodes a fast-twitch skeletal muscle protein, a member of the troponin I gene

family, and a component of the troponin complex including troponin T, troponin C and troponin I subunits. The troponin complex, along with tropomyosin, is responsible for the calcium-dependent regulation of striated muscle contraction. Mouse studies show that this component is also present in vascular smooth muscle and may play a role in regulation of smooth muscle function. In addition to muscle tissues, this protein is found in corneal epithelium, cartilage where it is an inhibitor of angiogenesis to inhibit tumor growth and metastasis, and mammary gland where it functions as a co-activator of estrogen receptor-related receptor alpha. This protein also suppresses tumor growth in human ovarian carcinoma. Mutations in this gene cause myopathy and distal arthrogryposis type 2B. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq,

Mar 2009]

Product images:



Coomassie blue staining of purified TNNI2 protein (Cat# [TP305676]). The protein was produced from HEK293T cells transfected with TNNI2 cDNA clone (Cat# [RC205676]) using MegaTran 2.0 (Cat# [TT210002]).