

## **Product datasheet for TP305689M**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## KIN (NM\_012311) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human KIN, antigenic determinant of recA protein homolog (mouse)

(KIN), 100 μg

Species: Human Expression Host: HEK293T

**Expression cDNA** >RC205689 protein sequence **Clone or AA Sequence:** Red=Cloning site Green=Tags(s)

MGKSDFLTPKAIANRIKSKGLQKLRWYCQMCQKQCRDENGFKCHCMSESHQRQLLLASENPQQFMDYFSE EFRNDFLELLRRRFGTKRVHNNIVYNEYISHREHIHMNATQWETLTDFTKWLGREGLCKVDETPKGWYIQ YIDRDPETIRRQLELEKKKKQDLDDEEKTAKFIEEQVRRGLEGKEQEVPTFTELSRENDEEKVTFNLSKG ACSSSGATSSKSSTLGPSALKTIGSSASVKRKESSQSSTQSKEKKKKKSALDEIMEIEEEKKRTARTDYW LQPEIIVKIITKKLGEKYHKKKAIVKEVIDKYTAVVKMIDSGDKLKLDQTHLETVIPAPGKRILVLNGGY

RGNEGTLESINEKTFSATIVIETGPLKGRRVEGIQYEDISKLA

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK
Predicted MW: 45.2 kDa

**Concentration:**  $>0.05 \mu g/\mu L$  as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling

conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 036443





**Locus ID:** 22944

UniProt ID: O60870

RefSeq Size: 6401

Cytogenetics: 10p14

RefSeq ORF: 1179


Synonyms: BTCD; KIN17; Rts2

**Summary:** The protein encoded by this gene is a nuclear protein that forms intranuclear foci during

proliferation and is redistributed in the nucleoplasm during the cell cycle. Short-wave ultraviolet light provokes the relocalization of the protein, suggesting its participation in the cellular response to DNA damage. Originally selected based on protein-binding with RecA antibodies, the mouse protein presents a limited similarity with a functional domain of the bacterial RecA protein, a characteristic shared by this human ortholog. Alternative splicing of this gene results

in multiple transcript variants. [provided by RefSeq, Jan 2012]

## **Product images:**



Coomassie blue staining of purified KIN protein (Cat# [TP305689]). The protein was produced from HEK293T cells transfected with KIN cDNA clone (Cat# [RC205689]) using MegaTran 2.0 (Cat# [TT210002]).