

## Product datasheet for TP305694M

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PCID1 (EIF3M) (NM 006360) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human eukaryotic translation initiation factor 3, subunit M (EIF3M), 100

μg

Species: Human Expression Host: HEK293T

**Expression cDNA Clone** >RC205694 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MSVPAFIDISEEDQAAELRAYLKSKGAEISEENSEGGLHVDLAQIIEACDVCLKEDDKDVESVMNSVVSL LLILEPDKQEALIESLCEKLVKFREGERPSLRLQLLSNLFHGMDKNTPVRYTVYCSLIKVAASCGAIQYI PTELDQVRKWISDWNLTTEKKHTLLRLLYEALVDCKKSDAASKVMVELLGSYTEDNASQARVDAHRCIVR ALKDPNAFLFDHLLTLKPVKFLEGELIHDLLTIFVSAKLASYVKFYQNNKDFIDSLGLLHEQNMAKMRLL TFMGMAVENKEISFDTMQQELQIGADDVEAFVIDAVRTKMVYCKIDQTQRKVVVSHSTHRTFGKQQWQQL

YDTLNAWKQNLNKVKNSLLSLSDT

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK
Predicted MW: 42.3 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 006351



Locus ID: 10480

UniProt ID: Q7L2H7 RefSeq Size: 1338 Cytogenetics: 11p13 RefSeq ORF: 1122


Synonyms: B5; GA17; hfl-B5; PCID1; TANGO7

**Summary:** This gene encodes a protein that is part of the eurkaryotic translation initiation factor 3

complete (eIF-3) required for protein synthesis. Elevated levels of the encoded protein are present in cancer cell lines. Inactivation of the encoded protein has been shown to interfere with translation of herpes virus mRNAs by preventing the association of mRNAs with the ribosomes. A pseudogene of this gene is located on the X chromosome. [provided by RefSeq,

Dec 2011]

## **Product images:**



Coomassie blue staining of purified EIF3M protein (Cat# [TP305694]). The protein was produced from HEK293T cells transfected with EIF3M cDNA clone (Cat# [RC205694]) using MegaTran 2.0