

Product datasheet for TP305697L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NTAN1 (NM_173474) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human N-terminal asparagine amidase (NTAN1), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC205697 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MPLLVEGRRVRLPQSAGDLVRAHPPLEERARLLRGQSVQQVGPQGLLYVQQRELAVTSPKDGSISILGSD DATTCHIVVLRHTGNGATCLTHCDGTDTKAEVPLIMNSIKSFSDHAQCGRLEVHLVGGFSDDRQLSQKLT HQLLSEFDRQEDDIHLVTLCVTELNDREENENHFPVIYGIAVNIKTAEIYRASFQDRGPEEQLRAARTLA GGPMISIYDAETEQLRIGPYSWTPFPHVDFWLHQDDKQILENLSTSPLAEPPHFVEHIRSTLMFLKKHPS

PAHTLFSGNKALLYKKNEDGLWEKISSPGS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 34.5 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 775745

Locus ID: 123803



NTAN1 (NM_173474) Human Recombinant Protein - TP305697L

UniProt ID: Q96AB6

RefSeq Size: 1241

Cytogenetics: 16p13.11

RefSeq ORF: 930


Synonyms: PNAA; PNAD

Summary: The protein encoded by this gene functions in a step-wise process of protein degradation

through the N-end rule pathway. This protein acts as a tertiary destabilizing enzyme that deamidates N-terminal L-Asn residues on proteins to produce N-terminal L-Asp. L-Asp substrates are subsequently conjugated to L-Arg, which is recognized by specific E3 ubiquitin ligases and targeted to the proteasome. Pseudogenes of this gene are located on the long arms of chromosomes 8, 10 and 12. Alternative splicing results in multiple transcript variants

that encode different protein isoforms. [provided by RefSeq, Jul 2012]

Product images:



Coomassie blue staining of purified NTAN1 protein (Cat# [TP305697]). The protein was produced from HEK293T cells transfected with NTAN1 cDNA clone (Cat# [RC205697]) using MegaTran 2.0 (Cat# [TT210002]).