

Product datasheet for TP305715

RPA34 (RPA2) (NM_002946) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human replication protein A2, 32kDa (RPA2), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC205715 protein sequence Red =Cloning site Green =Tags(s) MWNSGFESYGSSSYGGAGGYTQSPGGFGSPAPSQAEKKSRRARAHIVPCTISQLLSATLVDEVFRIGNVE ISQVTIVGIIRHAEKAPTNIIVYKIDDMTAAPMDVRQWVDTDTSSENTVVPETYVKVAGHLRSFQNKKS LVAFKIMPLEDMNEFTTHILEVINAHMVLSKANSQPSAGRAPISNPGMSEAGNFGGNSFMPANGLTVAQ N QVLNLIKACPRPEGLNFQDLKNQLKHMSVSSIKQAVDFLSNEGHIYSTVDDDDHFKSTDAE SGPTRTRPLE QKLISEEDLAANDILDYKDDDDK V
Tag:	C-Myc/DDK
Predicted MW:	29.1 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_002937</u>
Locus ID:	6118


[View online »](#)

UniProt ID: [P15927](#)

RefSeq Size: 1819

Cytogenetics: 1p35.3

RefSeq ORF: 810


Synonyms: REPA2; RP-A p32; RP-A p34; RPA32

Summary: This gene encodes a subunit of the heterotrimeric Replication Protein A (RPA) complex, which binds to single-stranded DNA (ssDNA), forming a nucleoprotein complex that plays an important role in DNA metabolism, being involved in DNA replication, repair, recombination, telomere maintenance, and co-ordinating the cellular response to DNA damage through activation of the ataxia telangiectasia and Rad3-related protein (ATR) kinase. The RPA complex protects single-stranded DNA from nucleases, prevents formation of secondary structures that would interfere with repair, and co-ordinates the recruitment and departure of different genome maintenance factors. The heterotrimeric complex has two different modes of ssDNA binding, a low-affinity and high-affinity mode, determined by which oligonucleotide/oligosaccharide-binding (OB) domains of the complex are utilized, and differing in the length of DNA bound. This subunit contains a single OB domain that participates in high-affinity DNA binding and also contains a winged helix domain at its carboxy terminus, which interacts with many genome maintenance protein. Post-translational modifications of the RPA complex also plays a role in co-ordinating different damage response pathways. [provided by RefSeq, Sep 2017]

Protein Families: Druggable Genome, Stem cell - Pluripotency

Protein Pathways: DNA replication, Homologous recombination, Mismatch repair, Nucleotide excision repair

Product images:



Coomassie blue staining of purified RPA2 protein (Cat# TP305715). The protein was produced from HEK293T cells transfected with RPA2 cDNA clone (Cat# [RC205715]) using MegaTran 2.0 (Cat# [TT210002]).