

Product datasheet for TP305718L

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RRM2 (NM_001034) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human ribonucleotide reductase M2 polypeptide (RRM2), 1 mg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC205718 representing NM_001034

or AA Sequence: Red=Cloning site Green=Tags(s)

MLSLRVPLAPITDPQQLQLSPLKGLSLVDKENTPPALSGTRVLASKTARRIFQEPTEPKTKAAAPGVEDE PLLRENPRRFVIFPIEYHDIWQMYKKAEASFWTAEEVDLSKDIQHWESLKPEERYFISHVLAFFAASDGI VNENLVERFSQEVQITEARCFYGFQIAMENIHSEMYSLLIDTYIKDPKEREFLFNAIETMPCVKKKADWA LRWIGDKEATYGERVVAFAAVEGIFFSGSFASIFWLKKRGLMPGLTFSNELISRDEGLHCDFACLMFKHL VHKPSEERVREIIINAVRIEQEFLTEALPVKLIGMNCTLMKQYIEFVADRLMLELGFSKVFRVENPFDFM

ENISLEGKTNFFEKRVGEYQRMGVMSSPTENSFTLDADF

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 44.7 kDa

Concentration: $>0.05 \mu g/\mu L$ as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Bioactivity: Binding assay (PMID: <u>29765556</u>)

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 001025



Locus ID: 6241

UniProt ID: P31350
RefSeq Size: 2500
Cytogenetics: 2p25.1
RefSeq ORF: 1167

Synonyms: C2orf48; R2; RR2; RR2M

Summary: This gene encodes one of two non-identical subunits for ribonucleotide reductase. This

reductase catalyzes the formation of deoxyribonucleotides from ribonucleotides. Synthesis of the encoded protein (M2) is regulated in a cell-cycle dependent fashion. Transcription from this gene can initiate from alternative promoters, which results in two isoforms that differ in the lengths of their N-termini. Related pseudogenes have been identified on chromosomes 1


and X. [provided by RefSeq, Sep 2009]

Protein Families: Druggable Genome

Protein Pathways: Glutathione metabolism, Metabolic pathways, p53 signaling pathway, Purine metabolism,

Pyrimidine metabolism

Product images:



Coomassie blue staining of purified RRM2 protein (Cat# [TP305718]). The protein was produced from HEK293T cells transfected with RRM2 cDNA clone (Cat# [RC205718]) using MegaTran 2.0 (Cat# [TT210002]).