

Product datasheet for TP305723

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Proteasome subunit beta type 4 (PSMB4) (NM 002796) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human proteasome (prosome, macropain) subunit, beta type, 4

(PSMB4), 20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC205723 representing NM_002796

or AA Sequence: Red=Cloning site Green=Tags(s)

MEAFLGSRSGLWAGGPAPGQFYRIPSTPDSFMDPASALYRGPITRTQNPMVTGTSVLGVKFEGGVVIAAD MLGSYGSLARFRNISRIMRVNNSTMLGASGDYADFQYLKQVLGQMVIDEELLGDGHSYSPRAIHSWLTRA MYSRRSKMNPLWNTMVIGGYADGESFLGYVDMLGVAYEAPSLATGYGAYLAQPLLREVLEKQPVLSQTEA

RDLVERCMRVLYYRDARSYNRFQIATVTEKGVEIEGPLSTETNWDIAHMISGFE

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 29 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 002787

Locus ID: 5692





UniProt ID: P28070, A0A140VK46

925 RefSeq Size:

Cytogenetics: 1q21.3 792 RefSeq ORF:

HN3; HsN3; PRAAS3; PROS-26; PROS26 Synonyms:

Summary: The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S


core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that

is a 20S core beta subunit. [provided by RefSeg, Jul 2008]

Protein Families: Druggable Genome, Protease

Protein Pathways: Proteasome

Product images:



Coomassie blue staining of purified PSMB4 protein (Cat# TP305723). The protein was produced from HEK293T cells transfected with PSMB4 cDNA clone (Cat# [RC205723]) using

MegaTran 2.0 (Cat# [TT210002]).