

Product datasheet for TP305782M

KCTD11 (NM_001002914) Human Recombinant Protein

Product data:


Product Type:	Recombinant Proteins
Description:	Recombinant protein of human potassium channel tetramerisation domain containing 11 (KCTD11), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC205782 protein sequence Red=Cloning site Green=Tags(s)
	MLGAMFRAGTPMPPNLNSQGGGHYFIDRDGKAFRHILNFLRLGRDLPRGYGETALLRAEADFYQIRPLL DALRELEASQGTPAPTAALLHADVDVSPRLVHFSARRGPHHYELSSVQVDTFRANLFCTDSECLGALRAR FGVASGDRAEGSPHFHLEWAPRPVELPEVEYGRLGLQPLWTGGPGERREVVGTSPFLEEVLRVALEHGFR LDSVFPDPEDLLNSRSLRFVRH
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	25.7 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_001002914</u>
Locus ID:	147040


[View online »](#)

UniProt ID:	<u>Q693B1</u>
RefSeq Size:	3081
Cytogenetics:	17p13.1
RefSeq ORF:	696
Synonyms:	C17orf36; KCASH1; REN; REN/KCTD11
Summary:	Plays a role as a marker and a regulator of neuronal differentiation; Up-regulated by a variety of neurogenic signals, such as retinoic acid, epidermal growth factor/EGF and NGFB/nerve growth factor. Induces apoptosis, growth arrest and the expression of cyclin-dependent kinase inhibitor CDKN1B. Plays a role as a tumor repressor and inhibits cell growth and tumorigenicity of medulloblastoma (MDB). Acts as probable substrate-specific adapter for a BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex towards HDAC1. Functions as antagonist of the Hedgehog pathway on cell proliferation and differentiation by affecting the nuclear transfer of transcription factor GLI1, thus maintaining cerebellar granule cells in undifferentiated state, this effect probably occurs via HDAC1 down-regulation, keeping GLI1 acetylated and inactive. When knock-down, Hedgehog antagonism is impaired and proliferation of granule cells is sustained. Activates the caspase cascade.[UniProtKB/Swiss-Prot Function]

Protein Families: Ion Channels: Other

Product images:



Coomassie blue staining of purified KCTD11 protein (Cat# [TP305782]). The protein was produced from HEK293T cells transfected with KCTD11 cDNA clone (Cat# [RC205782]) using MegaTran 2.0 (Cat# [TT210002]).