

Product datasheet for TP305992M

KTEL1 (POGLUT1) (NM_152305) Human Recombinant Protein

Product data:


| | |
|---------------------------------------|--|
| Product Type: | Recombinant Proteins |
| Description: | Recombinant protein of human KTEL (Lys-Tyr-Glu-Leu) containing 1 (KTELC1), transcript variant 1, 100 µg |
| Species: | Human |
| Expression Host: | HEK293T |
| Expression cDNA Clone or AA Sequence: | >RC205992 protein sequence Red =Cloning site Green =Tags(s) |
| | MEWWASSPLRLWLLLFLLPSAQGRQKESGSKWKVFIDQINRSLENYEPCSSQNCSCYHGVIEEDLTPFRG GISRRMMAEVRRKLGTHYQITKNRLYREND CMFPSRCGVEHFILEVIGRLPDMEMVINVRDYPQVPKW MEPAIPVFSFKTSEYHDIMYPAWTFWEGGPAWPIYPTGLGRWDLFREDLVRSAQWPWKKNSTAYFR GSRTSPERDPLILLSRKNTKLVDAEYTKNQAWKSMKDTLGKPAKDVHLVDHCKYKYLNFNRGVAASFRF KHLFLCGSLVFHVGDEWLEFFYPQLKPWWHYIPVKTDLSNVQELLQFVKANDDVAQEIAERGSQFIRNHL QMDDITCYWENLLSEYSKFLSYNVTRRKGYDQIIPKMLKTEL |
| | TRTRPLEQKLISEEDLAANDILDYKDDDDKV |
| Tag: | C-Myc/DDK |
| Predicted MW: | 46 kDa |
| Concentration: | >0.05 µg/µL as determined by microplate BCA method |
| Purity: | > 80% as determined by SDS-PAGE and Coomassie blue staining |
| Buffer: | 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol |
| Preparation: | Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps. |
| Note: | For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. |
| Storage: | Store at -80°C. |
| Stability: | Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles. |
| RefSeq: | <u>NP_689518</u> |



[View online »](#)

| | |
|---------------|--|
| Locus ID: | 56983 |
| UniProt ID: | Q8NBL1 |
| RefSeq Size: | 3552 |
| Cytogenetics: | 3q13.33 |
| RefSeq ORF: | 1176 |
| Synonyms: | C3orf9; CLP46; hCLP46; KDELCL1; KTELC1; LGMD2Z; LGMDR21; MDS010; MDSRP; Rumi |
| Summary: | This gene encodes a protein with both O-glucosyltransferase and O-xylosyltransferase activity which localizes to the lumen of the endoplasmic reticulum. This protein has a carboxy-terminal KTEL motif which is predicted to function as an endoplasmic reticulum retention signal. This gene is an essential regulator of Notch signalling and likely plays a role in cell fate and tissue formation during development. It may also play a role in the pathogenesis of leukemia. Mutations in this gene have been associated with the autosomal dominant genodermatosis Dowling-Degos disease 4. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr 2014] |

Product images:



Coomassie blue staining of purified POGLUT1 protein (Cat# [TP305992]). The protein was produced from HEK293T cells transfected with POGLUT1 cDNA clone (Cat# [RC205992]) using MegaTran 2.0 (Cat# [TT210002]).