

## Product datasheet for TP306101

### CCDC93 (NM\_019044) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human coiled-coil domain containing 93 (CCDC93), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC206101 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)

MGLPRGPEGQGLPEVETREDEEQNVKLTEILELLVAAGHFRARIKGLSPFDKVVGGMTWCITTCNFDVDV  
DLLFQENSTIGQKIALSEKIVSVLPRMKCPHQLEPHQIQGMDFIHIFPVVQWLVKRAIETKEEMGDYIRS  
YSVSQFQKTYSLPEDDDFIKRKEKAIKTVDLSEVYKPRRKYKRHQGAEELLDEESRIHATLLEYGRRYG  
FSCQSKMEKAEDKKTALPAGLSATEKADAHEEDELRAAEEQRIQSLMTKMTAMANEESRLTASSVGQIVG  
LCSAEIKQIVSEYAEKQSELSAEESPEKLGTSQLHRRKVISLNKQIAQKTKHLEELRASHTSLQARYNEA  
KKTLELTKTYSEKLDKEQAALIEKIESKADPSILQNLRALVAMNENLKSQEYEFKAHCREEMTRLQOEIEN  
LKAERAPRGDEKTLSSGEPGTLTSAMTHDEDLDRRYNMEKEKLYKIRLLQARRNREIAILHRKIDEVPS  
RAELIQYQKRFIELYRQISAVHKETKQFFTLNLTDDKKVYLEKEISLLNSIHENFQSAMASPAARDQFL  
RQMEQIVEGIKQSRMKMEKKKQENKMRRDQLNDQYLELLEKQRLYFKTVKEFKEEGRKNEMLLSKVKAKA  
S

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV**


Tag:	C-Myc/DDK
Predicted MW:	73 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.



[View online »](#)

<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<a href="#">NP_061917</a>
<b>Locus ID:</b>	54520
<b>UniProt ID:</b>	<a href="#">Q567U6</a>
<b>RefSeq Size:</b>	6946
<b>Cytogenetics:</b>	2q14.1
<b>RefSeq ORF:</b>	1893
<b>Summary:</b>	Component of the CCC complex, which is involved in the regulation of endosomal recycling of surface proteins, including integrins, signaling receptor and channels. The CCC complex associates with SNX17, retriever and WASH complexes to prevent lysosomal degradation and promote cell surface recycling of numerous cargos such as integrins ITGA5:ITGB1 (PubMed:28892079, PubMed:25355947). Involved in copper-dependent ATP7A trafficking between the trans-Golgi network and vesicles in the cell periphery; the function is proposed to depend on its association within the CCC complex and cooperation with the WASH complex on early endosomes and is dependent on its interaction with WASHC2C (PubMed:25355947). [UniProtKB/Swiss-Prot Function]

### Product images:



Coomassie blue staining of purified CCDC93 protein (Cat# TP306101). The protein was produced from HEK293T cells transfected with CCDC93 cDNA clone (Cat# [RC206101]) using MegaTran 2.0 (Cat# [TT210002]).