

## Product datasheet for **TP306188L**

### **AADACL1 (NCEH1) (NM\_020792) Human Recombinant Protein**

#### Product data:


<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	Recombinant protein of human arylacetamide deacetylase-like 1 (AADACL1), 1 mg
<b>Species:</b>	Human
<b>Expression Host:</b>	HEK293T
<b>Expression cDNA Clone or AA Sequence:</b>	>RC206188 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	MSSCRGQKVAGGLRVWSPFPLCQPAGEPSRGKMRSSCVLLTALVALAAYVYIPLPGSVSDPWKMLLLDA TFRGAQQVSNLIHYLGLSHHLLALNFIIVSFGKKS AWSSAQVKVTD TDFDGV E V R V F E G P P K P E E P L K R S VYIHGGGWALASAKIRYYDELCTAMAEELNAVIVSIEYRLVPKVFPEQIHDVVRATKYFLKPEVLQKY MVDPGRICISGDSAGGNLAAALGQQFTQDASLKNKLLQALIYPVLQALDFNTPSYQQNVNTPILPRYVM VKYWVDYFKGNYDFVQAMIVNNHTSLDVEEAAAVRARLNWTSLLPASFTKNYKPVVQTTGNARIVQELPQ LLDARSAPLIADQAVLQLLPKTYILTCEHDVLRDDGIMYAKRLESAGVEVTLDFHFDGFGHGMIFTSWPT NFSVGIRTRNSYIKWLDQNL
	<b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
<b>Tag:</b>	C-Myc/DDK
<b>Predicted MW:</b>	48.9 kDa
<b>Concentration:</b>	>0.05 µg/µL as determined by microplate BCA method
<b>Purity:</b>	> 80% as determined by SDS-PAGE and Coomassie blue staining
<b>Buffer:</b>	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
<b>Preparation:</b>	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
<b>Note:</b>	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
<b>Storage:</b>	Store at -80°C.
<b>Stability:</b>	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
<b>RefSeq:</b>	<u><a href="#">NP_065843</a></u>



[View online »](#)

Locus ID:	57552
UniProt ID:	<a href="#">Q6PIU2</a> , <a href="#">A0A0R4J2G3</a>
RefSeq Size:	4294
Cytogenetics:	3q26.31
RefSeq ORF:	1320
Synonyms:	AADACL1; NCEH
Summary:	Hydrolyzes 2-acetyl monoalkylglycerol ether, the penultimate precursor of the pathway for de novo synthesis of platelet-activating factor. May be responsible for cholesterol ester hydrolysis in macrophages, thereby contributing to the development of atherosclerosis. Also involved in organ detoxification by hydrolyzing exogenous organophosphorus compounds. May contribute to cancer pathogenesis by promoting tumor cell migration.[UniProtKB/Swiss-Prot Function]
Protein Families:	Transmembrane

### Product images:



Coomassie blue staining of purified NCEH1 protein (Cat# [TP306188]). The protein was produced from HEK293T cells transfected with NCEH1 cDNA clone (Cat# [RC206188]) using MegaTran 2.0 (Cat# [TT210002]).