

Product datasheet for TP306226

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CIPC (NM_033426) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human KIAA1737 (KIAA1737), 20 μg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC206226 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MERKNPSRESPRRLSAKVGKGTEMKKVARQFGMAAAESDKDSGFSDGSSECLSSAEQMESEDMLSALGWS REDRPRQNSKTAKNAFPTLSPMVVMKNVLVKQGSSSSQLQSWTVQPSFEVISAQPQLLFLHPPVPSPVSP CHTGEKKSDSRNYLPILNSYTKIAPHPGKRGLSLGPEEKGTSGVQKKICTERLGPSLSSSEPTKAGAVPS SPSTPAPPSAKLAEDSALQGVPSLVAGGSPQTLQPVSSSHVAKAPSLTFASPASPVCASDSTLHGLESNS PLSPLSANYSSPLWAAEHLCRSPDIFSEQRQSKHRRFQNTLVVLHKSGLLEITLKTKELIRQNQATQVEL

DQLKEQTQLFIEATKSRAPQAWAKLQASLTPGSSNTGSDLEAFSDHPAI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 42.5 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeg: NP 219494

Locus ID: 85457





UniProt ID: Q9C0C6

RefSeq Size: 4397

Cytogenetics: 14q24.3 RefSeq ORF: 1197


Synonyms: KIAA1737

Summary: Transcriptional repressor which may act as a negative-feedback regulator of CLOCK-

ARNTL/BMAL1 transcriptional activity in the circadian-clock mechanism. May stimulate ARNTL/BMAL1-dependent phosphorylation of CLOCK. However, the physiogical relevance of these observations is unsure, since experiments in an animal model showed that CIPC is not

critially required for basic circadian clock.[UniProtKB/Swiss-Prot Function]

Product images:



Coomassie blue staining of purified CIPC protein (Cat# TP306226). The protein was produced from HEK293T cells transfected with CIPC cDNA clone (Cat# [RC206226]) using MegaTran 2.0 (Cat# [TT210002]).