

Product datasheet for **TP306273**

HAAO (NM_012205) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human 3-hydroxyanthranilate 3,4-dioxygenase (HAAO), 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC206273 protein sequence Red =Cloning site Green =Tags(s)

MERRLGVRAWVKENRGSFQPPVCNKLHQEQQLKVMFIGGPNTRKDYHIEEGEEVFYQLEGDMVLRVLEQG
KHRDVIRQGEIFLLPARVPHSPQRFANTVGLVVERRRLETLDGLRYYVGDMDVLFKWFYCKDLGTQ
LAPIIQEFFSSEQYRTGKPIPDQLLKEPPFPLSTRSIMEPMSLDAWLDSHHRELQAGTPLSLFGDTYETQ
VIAYGQGSSEGLRQNVVWLWQLEGSSVVTMGGRRSLAPDSSLVLAGTSYAWERTQGSVALSVTQDPA
CKKPLG

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	32.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	NP_036337
Locus ID:	23498



[View online »](#)

UniProt ID: [P46952](#)

RefSeq Size: 1301

Cytogenetics: 2p21


RefSeq ORF: 858

Synonyms: 3-HAO; h3HAO; HAO; VCRL1

Summary: 3-Hydroxyanthranilate 3,4-dioxygenase is a monomeric cytosolic protein belonging to the family of intramolecular dioxygenases containing nonheme ferrous iron. It is widely distributed in peripheral organs, such as liver and kidney, and is also present in low amounts in the central nervous system. HAAO catalyzes the synthesis of quinolinic acid (QUIN) from 3-hydroxyanthranilic acid. QUIN is an excitotoxin whose toxicity is mediated by its ability to activate glutamate N-methyl-D-aspartate receptors. Increased cerebral levels of QUIN may participate in the pathogenesis of neurologic and inflammatory disorders. HAAO has been suggested to play a role in disorders associated with altered tissue levels of QUIN. [provided by RefSeq, Jul 2008]

Protein Pathways: Metabolic pathways, Tryptophan metabolism

Product images:



Coomassie blue staining of purified HAAO protein (Cat# TP306273). The protein was produced from HEK293T cells transfected with HAAO cDNA clone (Cat# [RC206273]) using MegaTran 2.0 (Cat# [TT210002]).