

Product datasheet for TP306301

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

EIF5 (NM_001969) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human eukaryotic translation initiation factor 5 (EIF5), transcript

variant 1, 20 µg

Species: Human
Expression Host: HEK293T

Expression cDNA Clone >RC206301 protein sequence or AA Sequence: Red=Cloning site Green=Tags(s)

MSVNVNRSVSDQFYRYKMPRLIAKVEGKGNGIKTVIVNMVDVAKALNRPPTYPTKYFGCELGAQTQFDVK NDRYIVNGSHEANKLQDMLDGFIKKFVLCPECENPETDLHVNPKKQTIGNSCKACGYRGMLDTHHKLCTF ILKNPPENSDSGTGKKEKEKKNRKGKDKENGSVSSSETPPPPPPPNEINPPPHTMEEEEDDDWGEDTTEE AQRRMDEISDHAKVLTLSDDLERTIEERVNILFDFVKKKKEEGVIDSSDKEIVAEAERLDVKAMGPLVL TEVLFNEKIREQIKKYRRHFLRFCHNNKKAKRYLLHGLECVVAMHQAQLISKIPHILKEMYDADLLEEEV IISWSEKASKKYVSKELAKEIRVKAEPFIKWLKEAEEESSGGEEEDEDENIEVVYSKAASVPKVETVKSD

NKDDDIDIDAI

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Tag: C-Myc/DDK

Predicted MW: 49 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by

conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.





RefSeq: NP 001960

Locus ID: 1983

UniProt ID: <u>P55010</u>, <u>A0A024R6Q1</u>

RefSeq Size: 5963

Cytogenetics: 14q32.32

RefSeg ORF: 1293


Synonyms: EIF-5; EIF-5A

Summary: Eukaryotic translation initiation factor-5 (EIF5) interacts with the 40S initiation complex to

promote hydrolysis of bound GTP with concomitant joining of the 60S ribosomal subunit to the 40S initiation complex. The resulting functional 80S ribosomal initiation complex is then active in peptidyl transfer and chain elongations (summary by Si et al., 1996 [PubMed

8663286]).[supplied by OMIM, May 2010]

Product images:



Coomassie blue staining of purified EIF5 protein (Cat# TP306301). The protein was produced from HEK293T cells transfected with EIF5 cDNA clone (Cat# [RC206301]) using MegaTran 2.0 (Cat# [TT210002]).