

Product datasheet for **TP306450M**

WFDC1 (NM_021197) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens WAP four-disulfide core domain 1 (WFDC1), 100 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA	>RC206450 representing NM_021197
Clone or AA Sequence:	Red=Cloning site Green=Tags(s)

MPLTGVGPGSCRRQIIRALCLLLLLHAGSAKNIWKRALPARLAEKSRAEEAGAPGGPRQPRADRCPPPP
RTLPPGACQAARCQADSECPRHRRCCYNGCAYACLEAVPPPPVLDWLVQPKPRWLGGNGWLLDGPPEVLQ
AEACSTTEDGAPELLCPSGYECHILSPGDVAEGIPNRGQCVKQRRQADGRILRHKLYKEYPEGDSKNVAE
PGRGQQKHFQ

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	20.6 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<u>NP_067020</u>
Locus ID:	58189



[View online »](#)

UniProt ID: [Q9HC57](#)

RefSeq Size: 1396

Cytogenetics: 16q24.1


RefSeq ORF: 660

Synonyms: PS20

Summary: This gene encodes a member of the WAP-type four disulfide core domain family. The WAP-type four-disulfide core domain contains eight cysteines forming four disulfide bonds at the core of the protein, and functions as a protease inhibitor in many family members. This gene is mapped to chromosome 16q24, an area of frequent loss of heterozygosity in cancers, including prostate, breast and hepatocellular cancers and Wilms' tumor. This gene is downregulated in many cancer types and may be involved in the inhibition of cell proliferation. The encoded protein may also play a role in the susceptibility of certain CD4 memory T cells to human immunodeficiency virus infection. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2013]

Protein Families: Secreted Protein

Product images:



Coomassie blue staining of purified WFDC1 protein (Cat# [TP306450]). The protein was produced from HEK293T cells transfected with WFDC1 cDNA clone (Cat# [RC206450]) using MegaTran 2.0 (Cat# [TT210002]).