

Product datasheet for **TP306527**

HCAR2 (NM_177551) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human niacin receptor 1 (NIACR1), 20 µg

Species: Human

Expression Host: HEK293T

Expression cDNA Clone or AA Sequence: >RC206527 protein sequence
Red=Cloning site **Green**=Tags(s)

MNRHHLQDHFLEIDKKNCCVFRDDFIVKVLPPVLGLEFIFGLLGNLALWIFCFHLKSWKSSRIFLFNLA
VADFLLIICLPFLMDNYVRRWDWKFWDIPCRMLFMLAMNRQGSIIFLTVAVDYFRVWHPHALNKIS
NRTAAIISCLLWGITIGLTVHLLKKKMPIQNGGANLCSFSICHTFQWHEAMFLEFFLPLGIILFCSAR
IIVSLRQRQMDRHAKIKRAITFIMVVAIVFVICFLPSVVRIRIFWLLHTSGTQNCVYRSVDLAFFITL
SFTYMNSMLDPVYYFSSPSFPNFFSTLINRCLQRKMTGEPDNNRSTSVELTGDPNKTRGAPEALMANS
EPWSPSYLGPTSP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Predicted MW: 41.7 kDa

Concentration: >0.05 µg/µL as determined by microplate BCA method

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

Bioactivity: Surface Plasmon Resonance (SPR) (PMID:[29473951](https://pubmed.ncbi.nlm.nih.gov/29473951/))

Preparation: Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.

Note: For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.


RefSeq: [NP_808219](https://ncbi.nlm.nih.gov/nuccore/NP_808219)



[View online »](#)

Locus ID:	338442
UniProt ID:	Q8TDS4
RefSeq Size:	2082
Cytogenetics:	12q24.31
RefSeq ORF:	1089
Synonyms:	GPR109A; HCA2; HM74a; HM74b; NIACR1; Puma-g; PUMAG
Summary:	Acts as a high affinity receptor for both nicotinic acid (also known as niacin) and (D)-beta-hydroxybutyrate and mediates increased adiponectin secretion and decreased lipolysis through G(i)-protein-mediated inhibition of adenylyl cyclase. This pharmacological effect requires nicotinic acid doses that are much higher than those provided by a normal diet. Mediates nicotinic acid-induced apoptosis in mature neutrophils. Receptor activation by nicotinic acid results in reduced cAMP levels which may affect activity of cAMP-dependent protein kinase A and phosphorylation of target proteins, leading to neutrophil apoptosis. The rank order of potency for the displacement of nicotinic acid binding is 5-methyl pyrazole-3-carboxylic acid = pyridine-3-acetic acid > acifran > 5-methyl nicotinic acid = acipimox >> nicotinuric acid = nicotinamide.[UniProtKB/Swiss-Prot Function]
Protein Families:	Druggable Genome, GPCR, Transmembrane

Product images:



Coomassie blue staining of purified HCAR2 protein (Cat# TP306527). The protein was produced from HEK293T cells transfected with HCAR2 cDNA clone (Cat# [RC206527]) using MegaTran 2.0 (Cat# [TT210002]).